Chapter 19 Measures for Acute Stress Disorder and Posttraumatic Stress Disorder

Susan M. Orsillo

Note: Throughout this chapter, reference is made to the various DSM-IV criteria for PTSD (American Psychiatric Association, 1994). Criterion A-1 refers to the occurrence of a traumatic event. Criterion A-2 refers to the presence of intense fear, helplessness, and horror at the time of the traumatic event. Criterion B refers to the presence of symptoms involving reexperiencing the traumatic event. Criterion C refers to symptoms of persistent avoidance of stimuli related to the traumatic event and numbing of general responsiveness. Criterion D includes symptoms of increased arousal since experiencing the trauma. Criterion E refers to the duration of the disturbance (at least 1 month for PTSD). Criterion F refers to the presence of clinically significant distress or impairment.

ACCIDENT FEAR QUESTIONNAIRE (AFQ)

Original Citation

Kuch, K., Cox, B. J., & Direnfeld, D. M. (1995). A brief self-rating scale for PTSD after road vehicle accident. Journal of Anxiety Disorders, 9, 503-514.

Susan M. Orsillo • National Center for PTSD–Women's Health Sciences Division, Boston VA Healthcare System, and Boston University School of Medicine, Boston, Massachusetts 02130.

255

Purpose

To measure PTSD-related phobic avoidance following involvement in a motor vehicle accident.

Description

The AFQ is a self-report scale consisting of an MVA profile that includes 10 yes/no questions about the accident and related anxiety, and 10 phobic avoidance items (AFQ-PA) in which the respondent is asked to rate his or her avoidance on a nine-point scale ranging from 0 (would not avoid it) to 8 (would always avoid it). There are also two descriptive questions about the accident and one question that assesses interference from physical illness (e.g., back pain) using the same nine-point scale.

Administration and Scoring

The AFQ can be administered in 5 to 10 minutes. The AFQ-PA subscale can be scored by summing the 10 items. A cutoff score of 15 on the AFQ-PA may be used to detect the presence of PTSD and/or accident phobia (see below).

Psychometric Properties

Sample Scores and Norms. The mean scores for the AFQ-PA are available from a sample of 54 men and women seeking treatment for pain or some other somatic symptom following a motor vehicle accident (Kuch et al., 1995). Individuals diagnosed with PTSD obtained a mean score of 54.44 (SD = 11.36); those diagnosed with accident phobia had a mean score of 34.00 (SD = 14.77); those with neither disorder had a mean score of 14.66 (SD = 12.46). Lower means were obtained in a sample of 113 accident victims referred to a research study from a rehabilitation center (Asmundson, Cox, Larsen, Frombach, & Norton, 1999). The mean score of individuals diagnosed with PTSD and accident phobia combined was 20.9 (SD = 14.0) and the comparison group obtained a mean of 9.0 (SD = 9.6). The percentages of respondents by diagnosis who endorsed each item in the accident profile are found in Kuch et al. (1995) and Asmundson et al. (1999).

Reliability. Items making up the MVA profile subscale had fairly low internal consistency ($\alpha = .67$) likely reflecting the divergent nature of the items (Asmundson et al., 1999). In contrast, good internal consistency was demonstrated for the AFQ-PA subscale (αs .80 to .89; Asmundson et al., 1999; Kuch et al., 1995).

Validity. The convergent and discriminant validity of the AFQ-PA was evaluated across a number of measures (Asmundson et al., 1999). The AFQ-PA was moderately associated with measures of anxiety sensitivity (.43), alexithymia (.33), and somatization (.28). The measure was not associated with extraversion and perceived self-control.

Individuals diagnosed with PTSD scored significantly higher than those with accident phobia and those with neither disorder in one sample (Kuch et al., 1995). In another sample, the PTSD and phobia groups did not differ, but these groups combined scored significantly higher than individuals with neither disorder (Asmundson et al., 1999).

MEASURES FOR ACUTE STRESS DISORDER AND PTSD | 257

The diagnostic efficiency of the AFQ-PA was assessed against a structured clinical interview. A cutoff score of 15 on the AFQ-PA appeared to be the optimal score for screening. However, this score yielded a sensitivity of only .67 and specificity of .78 with 55% of those scoring at or above the cutoff meeting diagnostic criteria for PTSD or accident phobia and 85% of those below the cutoff not meeting criteria for either disorder.

Source

The AFQ is reprinted in the original citation and in Appendix B. More information can be obtained by contacting Klaus Kuch, M.D. Forensic Program (4th Floor), Centre for Addiction and Mental Health, Clarke Institute Division, 250 College Street, Toronto, ON M5T 1R8, Canada; (tel) 705-487-2324; (e-mail) klaus.kuch@sympatico.ca.

ACUTE STRESS DISORDER INTERVIEW (ASDI)

Original Citation

Bryant, R. A., Harvey, A. G., Dang, S. T., & Sackville, T. (1998). Assessing acute stress disorder: Psychometric properties of a structured clinical interview. *Psychological Assessment*, 10, 215–220.

Purpose

To diagnose acute stress disorder.

Description

The ASDI is a clinician-rated scale consisting of 19 items that relate to criterion B (dissociation, 5 items), criterion C (reexperiencing, 4 items), criterion D (avoidance, 4 items), and criterion E (arousal, 6 items). Each item is scored dichotomously as 0 (symptom absent) or 1 (symptom present). The ASDI also includes items that assess the objective and subjective experience of the traumatic event (criterion A, 3 items), the duration of each symptom (criterion F), and impairment (criterion G, 4 items).

Administration and Scoring

The ASDI can be administered in 10 minutes. Scoring is according to DSM-IV criteria for ASD (see above).

Psychometric Properties

The psychometric properties of the ASDI were evaluated in a multisample, multistudy paper (Bryant et al., 1998).

Sample Scores and Norms. Thirteen patients out of 56 (23%) patients who were admitted to a hospital following a traumatic event met criteria for ASD. Twenty-four patients out of 60 (40%) who were referred to a PTSD unit following a traumatic event were diagnosed with ASD.

Reliability. The internal consistency for the 19 symptom items of the ASDI was found to be excellent (r = .90) among a sample of 65 patients admitted to a hospital following a traumatic event. The individual symptom clusters were lower: *dissociation* (r = .67), *reexperiencing* (r = .67), *avoidance* (r = .69), and *arousal* (r = .76).

Two- to seven-day test-retest reliability was reported for a sample of 60 adults seeking treatment. Correlations for each of the symptom clusters ranged from .80 to .87. Further, 88% of those who were diagnosed at time 1 were also diagnosed at time 2 and 94% of participants who were not diagnosed at time 1 also did not receive a diagnosis at time 2.

Validity. The items of the ASDI were rated by experts on a five-point scale ranging from 1 (not at all) to 5 (extremely) on their relevance (M = 4.86, SD = 0.93), specificity (M = 4.44, SD = 0.43), and clarity (M = 4.51, SD = 0.27). The content validity of the ASDI was evaluated in a sample of 56 inpatients admitted to a hospital following the occurrence of a traumatic event. The ASDI cluster score for the *dissociation* symptoms was significantly correlated with a measure of dissociation (r = .35), the *reexperiencing* cluster correlated with a measure of a voidance (r = .83), and the *arousal* cluster correlated with a measure of state anxiety (r = .38).

Given the lack of a gold standard measure of ASD, the ASDI was validated against the diagnosis of expert clinicians. The sensitivity of the ASDI was 91% and the specificity was 93%. Kappa values were .75 for the overall diagnosis, .79 for the stressor, .65 for *dissociation*, .61 for *reexperiencing*, .73 for *avoidance*, and .41 for *arousal*.

Alternative Forms

A self-report version of the ASDI, the Acute Stress Disorder Scale (ASDS) has also been developed (Bryant, Moulds, & Guthrie, 2000). Patients diagnosed with ASD scored a mean of 65.11 (SD = 14.74) on the ASDS and patients without ASD scored a mean of 36.97 (SD = 19.54). The internal consistency and test-retest reliability for the ASDS was found to be very good to excellent for the total score and the individual symptom clusters. Convergent validity and predictive validity against the interview version have also been established. The ASDS can be obtained by contacting Dr. Bryant (see source information below) and is reprinted in the original citation (Bryant et al., 2000).

Source

The ASDI can be obtained by contacting Richard A. Bryant, Ph.D., School of Psychology, University of New South Wales, Sydney, NSW 2052, Australia; (e-mail) r.bryant@ unsw.edu.au.

CLINICIAN-ADMINISTERED PTSD SCALE (CAPS)

Original Citations

Blake, D. D., Weathers, F. W., Nagy, L. M., Kaloupek, D. G., Klauminzer, G., Charney, D. S., & Keane, T. M. (1990). A clinician rating scale for assessing current and lifetime PTSD: The CAPS-1. The Behavior Therapist, 13, 187–188.

Blake, D. D., Weathers, F. W., Nagy, L. M., Kaloupek, D. G., Gusman, F. D., Charney, D. S., & Keane, T. M. (1995). The development of a clinician-administered PTSD scale. *Journal of Traumatic Stress*, 8, 75–90.

Purpose

To diagnose and assess symptoms of PTSD.

Description

The CAPS is a clinician-rated scale. The most up-to-date version includes a checklist of potentially traumatizing events. After it is administered, up to three events are chosen (based on their severity or recency) and a description of the event and the respondent's emotional response at the time of the event are obtained to establish DSM-IV criterion A. These events are referred to in the subsequent questions. Seventeen items directly assess DSM-IV criteria B, C, and D. Each item is rated on a five-point scale to determine the frequency (for most items 0 = never to 4 = daily or almost every day) and intensity (0 = none to 4 = extreme, with additional specific behavioral descriptions to each item). Raters are also permitted to indicate whether they believe each rating is of questionable validity (e.g., whether the patient is over- or underreporting). Criterion E is established by two questions on onset and duration. Criterion F is established by three questions on distress and impairment in functioning. Three items require the interviewer to make global ratings on the validity of responses, severity of PTSD and, if applicable, improvement since the previous assessment. If criteria are met for PTSD, five items tapping into associated features are administered. This version of the CAPS is able to assess symptoms over the past week, past month, and lifetime.

Administration and Scoring

The CAPS can be administered in 45-60 minutes. A total score is obtained by summing the frequency and intensity scores for each of the 17 symptom items. The CAPS can also be used to obtain a dichotomous rating of the presence or absence of PTSD. The psychometric properties of nine scoring rules (e.g., a symptom is present if the frequency rating is at least 1 and the intensity rating is at least 2; a symptom is present if the severity of a symptom {frequency + intensity} is greater than or equal to 4) have been examined (Weathers, Ruscio, & Keane, 1999). These authors concluded that the appropriate scoring rule should be based on the purpose of the assessment (e.g., screening versus differential diagnosis). Thus, it is highly recommended that users of the CAPS obtain this article.

Psychometric Properties

Sample Scores and Norms. In a sample of motor vehicle accident and sexual assault victims, the mean score on the CAPS was 45.9 (SD = 29.1; Blanchard, Jones-Alexander, Buckley, & Forneris, 1996).

Reliability. Internal consistency for intensity of PTSD symptom criteria was examined in a sample of 25 veterans (Blake et al., 1990). Cronbach's alpha ranged from .73 to .85. Similar results were found with a larger sample (Weathers & Litz, 1994). Internal consistency was also high within a sample of older veterans (α s range from .87 to .95; Hyer, Summers, Boyd, Litaker, & Boudewyns, 1996). Internater reliability on the same interview (with both raters present) was established within a sample of seven veterans for criteria B, C, and D (rs range from .92 to .99 for frequency and intensity; Blake et al., 1990). Diagnostic agreement within the pairs was perfect.

A more conservative test of interrater reliability was conducted in a larger sample of veterans. Three rater pairs independently interviewed veterans on two occasions 2 to 3 days apart (Weathers & Litz, 1994). The correlations for symptom clusters and total scores ranged from .77 to .98. However, the use of questionable validity ratings for each item was not shown to be reliable (Weathers & Litz, 1994).

Validity. A confirmatory factor analysis was conducted on a sample of 524 serviceseeking male veterans to examine the relative fit of a number of models hypothesized to reflect the dimensionality of PTSD (King, Leskin, King, & Weathers, 1998). The model of best fit was a four-factor, first order solution containing correlated factors reflecting the reexperiencing, effortful avoidance, emotional numbing, and hyperarousal symptom clusters of PTSD.

Convergent validity of the CAPS was demonstrated for a sample of 25 veterans (Blake et al., 1990). The CAPS was significantly correlated with self-report measures of PTSD (rs range from .70 to .84) and combat exposure (r = .42; Blake et al., 1990). In a sample of motor vehicle accident and sexual assault victims, the CAPS was significantly associated with a self-report measure of PTSD (r = .93). Within a large service-seeking veteran sample, score on the CAPS was shown to be significantly associated with other self-report measures of PTSD (rs range from .77 to .91), depression (rs range from .69 to .74), and anxiety (rs range from .65 to .76; Weathers & Litz, 1994). In contrast, in the same sample, the CAPS was only weakly associated with a measure of antisocial personality (r = .14). Further, when the effects of possible response bias were controlled for, the correlations with measures of PTSD remained strong, whereas the correlations with associated features dropped substantially (Weathers & Litz, 1994). Finally, convergent validity of the symptom subscales of the CAPS was assessed in a large sample of service-seeking veterans (King et al., 1998). A measure of state anxiety was more strongly associated with symptoms of hyperarousal and reexperiencing than the numbing and avoidance clusters. In contrast, numbing and hyperarousal were the symptom clusters most highly associated with depression (King et al., 1998).

The diagnostic utility of nine scoring rules for the CAPS was examined against a diagnosis obtained by structured clinical interview in a sample of service-seeking veterans (Weathers et al., 1999). All nine rules yielded efficiencies ranging from .82 to .87.

Diagnosis based on the CAPS has been shown to be predictive of heart rate reactivity in response to a combat-related priming event (e.g., Litz, Orsillo, Kaloupek, & Weathers, 2000). Treatment sensitivity was demonstrated in a study of trauma management therapy with veterans (Frueh, Turner, Beidel, Mirabella, & Jones, 1996), and in an open trial of exposure therapy with a mixed group of trauma survivors (Thompson, Charlton, Kerry, & Lee, 1995).

Alternative Forms

The CAPS has been translated into several languages including French, Spanish, Japanese, and Russian. A modification of the CAPS for use with Afghan refugees (Pushto and Farsi {Dari} languages) has been demonstrated to have good internal consistency and interrater reliability (Malekzai et al., 1996). The Dutch version has also been psychometrically examined (Hovens, van der Ploeg, Klaarenbeek, Schreuder, & Rivero, 1994a). A computerized version of the CAPS with good validity and reliability has been developed (Neal, Busuttil, Herapath, & Strike, 1994). A child version of the CAPS is available through the National Center for PTSD website. More information on the child version is available by contacting Elana Newman, University of Tulsa, Psychology/Lorton Hall, 600 S. College Ave., Tulsa, OK 74104-3189, USA; (e-mail) newmane@centum.utulsa.edu.

Source

More information about the CAPS, including a request form to obtain a copy of the measure, is available at the National Center for PTSD website (www.ncptsd.org/treatment/ assessment/caps.html). In addition, interested readers can contact Carole A. Goguen, Psy.D. at the National Center for PTSD (116D) VA Medical Center & Regional Office Center, 215 North Main St., White River Junction, VT 05009, USA. The CAPS can be obtained through the National Center for PTSD at no cost. In addition, a version of the CAPS is under development with Western Psychological Services. For more information, contact Western Psychological Services, 12031 Wilshire Boulevard, Los Angeles, CA 90025, USA; (tel) 800-648-8857; (fax) 310-478-7838; (e-mail) custsvc@wpspublish.com; (website) www.wpspublish.com.

DAVIDSON TRAUMA SCALE (DTS)

Original Citation

Davidson, J. R. T., Book, S. W., Colket, J. T., Tupler, L. A., Roth, S., David, D., Hertzberg, M., Mellman, T., Beckham, J. C., Smith, R. D., Davison, R. M., Katz, R., & Feldman, M. E. (1997). Assessment of a new self-rating scale for posttraumatic stress disorder. *Psychological Medicine*, 27, 153-160.

Purpose

To assess symptoms of PTSD among individuals with a history of trauma exposure.

Description

The DTS is a self-report scale comprised of 17 items corresponding to each of the DSM-IV symptoms of PTSD. For each item, the respondent rates frequency and severity for the previous week on a five-point scale. The *reexperiencing* symptoms are tied to a specific traumatic event described by the respondent. However, the *numbing*, *withdrawal*, and *hyper*-

262 CHAPTER 19

arousal events are not specifically linked to the traumatic event (e.g., the respondent is not asked if these symptoms arose as a result of, or at the time of, the trauma). For frequency, the scale ranges from 0 (not at all) to 4 (every day). For severity, the range is from 0 (not at all) to 4 (extremely distressing).

Administration and Scoring

The DTS takes 10 minutes to administer. A total score can be derived by summing all of the items. Subscale scores can be computed separately for frequency and severity. Subscale scores can also be computed separately for each of the symptom clusters: reexperiencing, avoidance, and hyperarousal.

Psychometric Properties

Sample Scores and Norms. Respondents derived from studies on veterans and victims of a natural disaster with PTSD obtained a mean score of 62 (SD = 38.0) versus a mean of 15.5 (SD = 13.8) for respondents without PTSD (Davidson, et al., 1997).

Reliability. Internal consistency was reported for participants in studies on rape victims, veterans, and victims of a natural disaster (Davidson, Book, et al., 1997). Cronbach's alpha for the combined sample ranged from .97 to .99 for the frequency only, severity only, and total items. Similar internal consistency ratings were found in a sample of survivors of childhood sexual assault (Zlotnick, Davidson, Shea, & Pearlstein, 1996).

Two-week test-retest data were available for 21 participants in a multicenter drug trial for individuals with a variety of trauma histories who were rated with "no change" on an independently derived measure of their symptoms (Davidson, et al., 1997). The test-retest reliability on this subsample was .86.

Factor analysis of a sample of individuals with a history of combat, rape, or natural disaster yielded two factors (Davidson, et al., 1997). The first accounting for 20% of the variance was interpreted as a severity factor. The second accounted for a small amount of the variance and consisted mostly of positive loadings on the *intrusive* items and negative loadings on the *avoidance* and *numbing* items. A factor analysis on only respondents with PTSD yielded six factors, the largest being similar to the severity factor discussed above.

Convergent validity was assessed with the same sample (Davidson, et al., 1997). Individuals diagnosed with PTSD on the basis of a structured clinical interview scored significantly higher than those without PTSD. A score of 40 on the DTS was associated with a sensitivity of 69%, specificity of 95%, and efficiency of 83%. Within the sample of patients treated with an antidepressant, DTS scores were significantly different (in the predicted directions) for five categories of PTSD severity (minimal, subclinical, clinical, severe, and very severe) determined by a physician's rating (Davidson, et al., 1997). Within the rape victim and veteran samples, the DTS was significantly correlated with a self-report (r = .64) and an interview (r = .78) measure of PTSD. The DTS was also significantly correlated with measures of general psychological distress (rs ranging from .44 to .65), but not with a measure of extroversion (r = .04).

In a sample of childhood sexual abuse survivors, DTS scores were significantly correlated with an interview measure of PTSD (rs ranged from .57 to .72 for frequency and severity), a measure of dissociation (rs ranged from .51 to .59), and a measure of affect regulation (rs ranged from .49 to .53; Zlotnick et al., 1996).

MEASURES FOR ACUTE STRESS DISORDER AND PTSD 263

Sensitivity to treatment effects was evaluated by comparing the DTS total score in responders versus nonresponders in the antidepressant trial (Davidson, et al., 1997). Responders demonstrated a significant decrease in DTS score, whereas nonresponders did not. Controlling for pretreatment scores, women with a history of childhood sexual assault who received a group treatment demonstrated a nonsignificant trend toward scoring lower on the DTS than individuals in the waitlist control condition (Zlotnick et al., 1996).

Alternative Forms

The DTS is available in French-Canadian and Spanish. A four-item scale called the SPAN (Startle, Physiological arousal, Anger, and Numbness) has been developed as a brief diagnostic screening from the DTS. The psychometric properties of this scale are described in Meltzer-Brody, Churchill, and Davidson (1999). The SPAN is also available from Multi-Health Systems (see below).

Source

The DTS is available through Multi-Health Systems Inc., 908 Niagara Falls Blvd., North Tonawanda, NY 14120-2060, USA; (tel) 800-456-3003 (USA) or 800-268-6011 (Canada); (fax) 416-424-1736; (webpage) www.mhs.com. A kit that contains a manual and 25 scoring forms is available for \$47.50 US. More information can also be obtained by contacting Jonathan R. T. Davidson, M.D., Department of Psychiatry, Duke University Medical Center, Box 3812, Durham, NC 27710, USA; (tel) 919-684-2880; (fax) 919-684-8866; (e-mail) tolme@acpub.duke.edu.

DISSOCIATIVE EXPERIENCES SCALE (DES)

Original Citations

Bernstein, E. M., & Putnam, F. W. (1986). Development, reliability and validity of a dissociation scale. *Journal of Nervous and Mental Disease*, 174, 727-735.

Carlson, E. B., & Putnam, F. W. (1993). An update on the Dissociative Experiences Scale. Dissociation, 6, 16–27.

Purpose

To measure frequency of dissociative experiences.

Description

The DES has been used in over 400 published studies on a variety of populations. Over 35 studies have been conducted on the psychometric properties of the scale. While the DES appears to be a clinically useful measure of dissociative features, it lacks some clarity because

it does not specify a time period for reporting and because it measures experiences that are reflective of both normal and pathological dissociation. The DES consists of 28 items that describe dissociative experiences including experiences of amnesia, depersonalization, derealization, imaginative involvement, and absorption. Respondents are asked to indicate the frequency (not including drug- or alcohol-related experiences) of these experiences using a 100-point scale. The original version of the DES used a visual analogue response scale consisting of a 100 mm line numerically anchored on the end points. A revised version uses a format of numbers from 0 to 100 (by 10s) and asks the respondent to circle the percentage of time that best reflects how much he or she has each experience in their daily life (no specific time window is indicated).

Administration and Scoring

The DES can be administered in 10 to 15 minutes. On the original version, the scale is scored by measuring the mark made by the respondent to the nearest 5 millimeters. On the revised version, the circled numbers are used. A total score is calculated by adding all of the items and dividing by 28. Subscale means based on a factor analysis (Carlson et al., 1991) can also be obtained, although, there is some debate over the validity of these factors (see psychometric review below). Amnestic dissociation is measured by taking the mean of items 3-6, 8, 10, 25, and 26. Absorption and imaginative involvement is measured by taking the mean of items 2, 14-18, 20, 22, and 23. Depersonalization and derealization is the mean of items 7, 11-13, 27, and 28. A score of 30 is used as a cutoff point for defining a respondent as high in dissociation.

Psychometric Properties

Sample Scores and Norms. Based on a meta-analysis of 85 studies on almost 12,000 respondents, means were derived for individuals with PTSD (N = 259, M = 32.58), for normal individuals (N = 1578, M = 11.57), and students (N = 5676, M = 14.27; van Ijzendoorn & Schuengel, 1996).

Reliability. The meta-analysis conducted by van Ijzendoorn and Schuengel (1996) included 16 studies that examined internal consistency, and the mean alpha across these studies was .93. The test-retest reliability of the DES has been shown to range from .79 to .96 over 4- to 8-week intervals (e.g., Bernstein & Putnam, 1986; Frischholz et al., 1990; Pitblado & Sanders, 1991).

Validity. Fischer and Elnitsky (1990) conducted a factor analysis on the DES in a student sample to determine if the three-factor solution (yielding the three subscales discussed earlier) would emerge. However, they found that a one-factor solution best accounted for the data. In contrast, Ross, Joshi, and Currie (1991) conducted a factor analysis on data derived from a large community sample and obtained the three-factor solution that was hypothesized. Carlson et al. (1991) conducted a factor analysis on DES scores derived from a multicenter study including 1574 individuals with and without a variety of psychological disorders (Carlson et al., 1991) and also obtained the three-factor solution. However, Waller (1995) reanalyzed their data set correcting for skewness and confirmed that one general factor best accounted for the variance.

The meta-analysis conducted by van Ijzendoorn and Schuengel (1996) included 26 studies that allowed for an examination of the convergent validity of the DES. The DES showed excellent convergent validity with other self-report and interview measures of dissociation (combined effect size d = 1.05, N = 1705), PTSD (combined effect size d = 0.75, N = 1099), and physical or sexual abuse (combined effect size d = 0.52, N = 2108).

In contrast, the discriminant validity of the DES is somewhat less well established. DES scores do not seem to be strongly associated with gender (combined effect size d = -0.01, N = 4074) or age (d = -0.24, N = 2474). However, DES scores have been shown to be significantly related to a number of measures of general psychological distress (rs ranging from .67 to .69; e.g., Walker, Katon, Neraas, Jemelka, & Massoth, 1992; Zlotnick et al., 1995). Although this may indicate that the DES measures general distress, it may also accurately reflect the relationship between general distress level and severity of dissociation.

Alternative Forms

The DES has been translated into at least 17 languages. There is an adolescent DES that is similarly formatted, but that contains different content, and is available through the Sidran Foundation (see below). A brief version of the DES, the DES-T, has been developed to specifically measure pathological dissociation (Waller, Putnam, & Carlson, 1996; Waller & Ross, 1997). A computerized version of the scale is available from Grant Fair, M.S.W., R.S.W., (e-mail) grantf@provcomm.net.

Source

The line version of the DES is reprinted in the Bernstein and Putnam (1986) paper, although the response line is not the correct length and one item is missing. The percentage version of the DES is reprinted in Carlson and Putnam (1993) and in Appendix B. One copy of the instrument (designate the language), a user's manual (the Carlson & Putnam, 1993, article referenced above), and a list of 333 references are available for purchase through the Sidran Foundation, 2328 West Joppa Road, Luterville, MD 21093, USA; (tel) 410-825-8888; (fax) 410-337-0747; (e-mail) sidran@sidran.org; (webpage) www.sidran.org.

DISTRESSING EVENT QUESTIONNAIRE (DEQ)

Original Citation

Kubany, E. S., Leisen, M. B., Kaplan, A. S., & Kelly, M. P. (2000). Validation of a brief measure of posttraumatic stress disorder: The Distressing Event Questionnaire (DEQ). *Psychological Assessment*, 12, 197-209.

Purpose

To assess PTSD and PTSD severity.

Description

The DEQ is a comprehensive measure that has been demonstrated to assess PTSD across a variety of trauma populations. Although the measure does not assess criterion A-1 (the occurrence of a traumatic event), it does assess criterion A-2, with three questions that assess the presence of intense fear, helplessness, and horror at the time of the traumatic event. The DEQ also includes 17 items that assess the diagnostic symptoms of PTSD (criteria B through D). Respondents are asked to indicate the degree to which they experienced each of the symptoms within the last month on a five-point scale ranging from 0 (absent or did not occur) to 4 (present to an extreme or severe degree). Criterion E is assessed by three questions that ask if the respondent had PTSD for more than 30 days, when the symptoms began, and how long they have persisted. Criterion F is assessed by 11 items that measure distress and impairment in various areas of functioning. Additional features associated with PTSD, including traumarelated guilt, anger, and unresolved grief over trauma-related losses, are also assessed.

Administration and Scoring

The DEQ can be administered in 5 to 7 minutes. A PTSD diagnosis can be obtained by following the DSM-IV diagnostic criteria or a symptom severity index can be obtained by summing the appropriate items. Cutoff scores differ for men and women (see below).

Psychometric Properties

Psychometric properties were evaluated in a multisample, multistudy study (Kubany et al., 2000).

Sample Scores and Norms. None are available.

Reliability. Reliability was assessed in a sample of male veterans. The internal consistency of the DEQ was excellent for the total score ($\alpha = .93$) and very good to excellent for each of the B, C, and D symptom clusters ($\alpha s .88$ to .91). Similar results were found for women with histories of sexual assault, abuse by an intimate partner, prostitution, and substance abuse.

Test-retest reliability for a male veteran sample (M = 17.5 days, SD = 12.3 days) for the overall scale was .95 with reliability coefficients (rs) for the various symptom clusters ranging from .69 to .72. Similar results were found among battered women (test-retest reliability for total score, r = .83; for subscales rs range from .76 to .81). Temporal stability of the DEQ for identifying the presence of a PTSD diagnosis was also demonstrated with the battered women. Utilizing all six DSM-IV criteria to establish a diagnosis resulted in 83% diagnostic agreement.

Validity. Six clinicians who specialize in PTSD rated relevance and representativeness of several aspects of the DEQ (e.g., response format, individual items) for measuring PTSD as defined in DSM-IV. The responses averaged "very well" to "considerably" relevant and representative for all indices.

Convergent validity was assessed in a male veteran sample. A sum of the 20 symptom items of the DEQ was significantly, positively correlated with another measure of PTSD (r = .83), a measure of depression (r = .76), and hostility (r = .55), and was negatively correlated with a measure of self-esteem (r = -.67).

MEASURES FOR ACUTE STRESS DISORDER AND PISD | 267

In a mixed sample of veterans and women with histories of sexual assault, abuse by an intimate partner, prostitution, and substance abuse, the DEQ was correlated with two other measures of PTSD (rs range from .82 to .94) and these correlations remained high across a variety of ethnic groups. Further, the sum of the additional items of the DEQ measuring associated features were also found to correlate significantly with three measures of PTSD, a measure of depression and a measure of guilt (rs = .71 to .91 for men and rs = .57 to .78 for women). In contrast, the DEQ was uncorrelated with a measure of social desirability in all samples except for within the group of women with a history of prostitution.

The ability of the DEQ to predict diagnostic status as assessed by another self-report measure of PTSD was examined. The percentage of diagnostic agreement between the two scales was 82%. PTSD designation on the DEQ was based on whether DSM-IV criteria B, C, and D were met using a symptom score of 2 (i.e., present to a moderate degree) or higher. The two measures agreed on positive PTSD cases 75% of the time, and agreement regarding the absence of PTSD occurred in 92% of the cases.

The discriminative validity of the DEQ was evaluated against the CAPS. A cutoff score of 26 for a veteran group correctly classified 86% of the sample. A cutoff score of 18 for a group of treatment-seeking women with histories of rape, incest, partner abuse, prostitution, and substance abuse correctly classified 90% of the sample.

Alternative Forms

There are different initial instruction versions of the DEQ depending on the purpose of the assessment and the setting in which it is conducted. Translations are available in Japanese and Tagalog. A computerized version of the scale (which incorporates this scale and the Traumatic Life Events Questionnaire) is currently being validated in a grant-funded study.

Source

The DEQ is available from Edward S. Kubany, Ph.D., National Center for PTSD, Department of Veterans Affairs, 1132 Bishop Street, Suite 307, Honolulu, HI 96813, USA; (tel) 808-566-1651; (fax) 808-566-1885; (e-mail) kubany@pixi.com. A published version of the DEQ is in development with Western Psychological Services, 12031 Wilshire Boulevard, Los Angeles, CA 90025, USA; (tel) 800-648-8857; (fax) 310-478-7838; (e-mail) custsvc@ wpspublish.com; (website) www.wpspublish.com.

IMPACT OF EVENT SCALE (IES)

Original Citations

Horowitz, M., Wilner, N., & Alvarez, W. (1979). Impact of Event Scale: A measure of subjective stress. *Psychosomatic Medicine*, 41, 209–218.

Weiss, D. S., & Marmar, C. R. (1997). The Impact of Event Scale-Revised. In J. P. Wilson & T. M. Keane (Eds.), Assessing psychological trauma and PTSD (pp. 399-411). New York: Guilford.

Purpose

To measure intrusion and avoidance resulting from exposure to traumatic events.

Description

The IES is a 15-item self-report questionnaire based on Horowitz's (1976) conceptualization of the stress response as including alternating phases of intrusions and avoidance. There are two subscales: *intrusion* and *avoidance* (see discussion below). Respondents are asked to indicate a specific life event and to rate the descriptive statements in response to that event. Respondents rate how frequently they have experienced each of the symptoms during the previous 7 days on a four-point scale. Weighted numerical ratings are assigned to the descriptors (not at all = 0, rarely = 1, sometimes = 3, and often = 5). Item values of 2 and 4 are not used. Although the IES can be used to assess responses to any type of stressful event, it has been widely used to measure symptoms of PTSD. However, the IES does not assess criterion D, hyperarousal symptoms (e.g., difficulty concentrating, exaggerated startle response). A revised version of the IES, discussed below, includes six items to assess this symptom cluster.

Administration and Scoring

It takes 10 minutes to administer the IES. The IES is scored by summing all of the items. Subscale scores can also be derived by summing the items that reflect *intrusion* (items 1, 4-6, 10, 11, 14) and *arousal* (items 2, 3, 7-9, 12, 13, 15). Horowitz (1982) identified total score thresholds for clinical concern as low (< 8.5), medium (8.6-19), and high (> 19). However, these cutoff points are not related to diagnostic status and their utility has been questioned (e.g., Joseph, 2000).

Psychometric Properties

Sample Scores and Norms. There are no published norms on the IES, but some individual studies report mean scores for different groups. For instance, the mean score on the IES in a sample of 130 service seeking veterans was 55.7 (SD = 10.6) for the total score and 27.6 (SD = 6.8) for the *intrusion* subscale and 28.2 (SD = 6.2) for the *avoidance* subscale (McFall, Smith, Roszell, Tarver, & Malas, 1990). Among a sample of survivors of a ferry disaster, the mean score was 35 for total score, 19 for *intrusion*, and 16 for *avoidance* (Joseph, Yule, Williams, & Hodgkinson, 1993). Female bank staff following an armed raid scored an average of 22.57 on the IES total score (Hodgkinson & Joseph, 1995).

Reliability. Good internal consistency for the total and subscale scores on the IES has been demonstrated in a sample of psychotherapy outpatients who had experienced a serious life event (α s range from .78 to .86; Horowitz et al., 1979) and a sample including outpatients and controls who had experienced parental bereavement (total score IES $\alpha = .86$; Zilberg, Weiss, & Horowitz, 1982).

Test-retest reliability over 1 week for a small sample of students who had recently dissected a cadaver was .87 for the total score, .89 for intrusion, and .79 for avoidance (Horo-

witz et al., 1979). Despite the fact that the IES is widely used, no other studies have reported test-retest reliability data on the scale, raising the concern that these findings might overestimate the true reliability of the scale (Joseph, 2000).

Validity. Psychotherapy outpatients reflecting on significantly distressing life events scored significantly higher on the *intrusion subscale*, *avoidance subscale*, and the total IES than medical students relating to their first dissection experience, which had occurred during the previous week (Horowitz et al., 1979). Patients seeking therapy for parental bereavement scored significantly higher than individuals recruited from the community who had also lost a parent (Zilberg et al., 1982).

Several factor analyses have confirmed the existence of the two hypothesized factors of the IES (e.g., Hodgkinson & Joseph, 1995; Joseph et al., 1993; Schwarzwald, Solomon, Weisenberg, & Mikulincer, 1987; Zilberg et al., 1982) with minor differences in the item loadings. However, even when the two-factor structure was supported, some of these studies have yielded a weaker, third factor that may in fact reflect the distinction of emotional avoidance or denial from active behavioral avoidance (e.g., Joseph et al., 1993; McDonald, 1997; Schwarzwald et al., 1987). Hodgkinson and Joseph (1995) found a change in the structure of a factor analysis over time. Specifically, they found *intrusion* to be a larger factor immediately posttrauma and *avoidance* to account for more variance at a later follow-up point (Hodgkinson & Joseph, 1995). Finally, other studies with more chronic trauma populations have reported a single factor solution, suggesting that the distinctiveness of intrusion and avoidance lessens over time (Hendrix, Jurich, & Schumm, 1994).

Studies have examined correlations between the subscales of the IES and have, for the most part, supported the notion that they measure separate but related constructs (rs = .40 to .78; Hodgkinson & Joseph, 1995; Horowitz et al., 1979, Neal, Busuttil, Rollins, et al., 1994; Zilberg et al., 1982). There is some evidence that the relationship between the two subscales changes over time (e.g., Zilberg et al., 1982).

The IES has been shown to be associated with another self-report measure of PTSD (rs for the total score and subscales ranged from .44 to .67) and a measure of general distress (rs range from .50 to .60) among women receiving inpatient treatment for trauma-related disorders (Allen, Coyne, & Huntoon, 1998). In a mixed sample of veterans seeking treatment and civilians, the IES was significantly associated with a self-report (rs ranged from .73 to .79) and a structured clinical interview measure of PTSD (rs ranged from .75 to .81; Neal, Busuttil, Rollins, et al., 1994). Further, the IES was significantly associated with general distress (rs ranged from .44 to .63) in a sample of women exposed to an armed raid (Hodgkinson & Joseph, 1995).

Neal, Busuttil, Rollins, et al. (1994) found that an optimum cutoff score of 35 for the total IES yielded a sensitivity of 89%, specificity of 88%, and overall diagnostic efficiency of 88%, relative to a structured clinical interview.

Treatment sensitivity has been demonstrated in several studies (Davidson et al., 1993; Foa, Rothbaum, Riggs, & Murdock, 1991; Horowitz et al., 1979).

The predictive validity of the IES has also been demonstrated. Perry, Difede, Musngi, Frances, and Jacobsberg (1992) found that IES *intrusion* scores 2 months posttrauma significantly predicted PTSD at 6 months and that IES *avoidance* scores at 6 months significantly predicted PTSD at 12 months posttrauma. Shalev, Peri, Canetti, and Schreiber (1996) found that IES scores 1 week posttrauma predicted PTSD 6 months later with 92.3% sensitivity, but only 34.2% specificity.

270 CHAPTER 19

Alternative Forms

The IES has been translated into several languages including Hebrew and Dutch. A revised version of the IES, which includes seven new items presumed to measure hyperarousal and one that measures flashback-type experiences, has been demonstrated to have very good internal consistency and moderate to good test-retest reliability (Weiss & Marmar, 1997). After collecting these psychometric data, the authors made some additional changes to the IES-R with regard to the instructions and the rating scale, which need to be empirically evaluated.

Source

The IES is reprinted in the original citation and in Appendix B. Additional information can be obtained by contacting Mardi Horowitz, M.D., University of California-San Francisco, P.O. Box 0984, Box F-LPP 357, San Francisco, CA 94142, USA. The IES-R is reprinted in Weiss and Marmar (1997). More information can be obtained by contacting Daniel S. Weiss, Ph.D., Director of PTSD Research, SFVAMC, Department of Psychiatry, University of California-San Francisco, San Francisco, CA 94143, USA; (e-mail) dweiss@itsa.ucsf.edu.

LOS ANGELES SYMPTOM CHECKLIST (LASC)

Original Citation

King, L. A., King, D. W., Leskin, G., & Foy, D. W. (1995). The Los Angeles Symptom Checklist: A self-report measure of posttraumatic stress disorder. *Assessment*, 2, 1–17.

Purpose

To measure PTSD symptoms from DSM-IV criteria B, C, and D as well as associated features.

Description

The LASC is a 43-item self-report scale. Seventeen of the items correspond fairly closely with the B, C, and D symptoms of PTSD. Each item is a word or phrase that is rated on a fivepoint scale ranging from 0 (not a problem) to 4 (extreme problem), reflecting the extent to which the symptom is a problem for the respondent. No time frame is established for rating symptoms.

Administration and Scoring

The LASC can be administered in 10 to 15 minutes. It can be scored in several ways. To be considered a positive PTSD case, a respondent must endorse an appropriate combination of symptoms with a rating of two or higher. Following DSM-IV criteria, the diagnosis of PTSD

MEASURES FOR ACUTE STRESS DISORDER AND PTSD | 271

requires at least one *reexperiencing* (B) symptom (items 5, 23, 28), three *avoidance* (C) symptoms (items 19, 29, 40-43), and two *arousal* (D) symptoms (items 1, 4, 8, 20, 25, 34, 37, 38). A partial PTSD diagnosis may be considered if a respondent endorses two of the three criteria. The LASC may also be scored as a continuous measure of PTSD severity, which requires summing the scores of the 17 items reflecting PTSD symptoms. Finally, the sum of all 43 items provides a global assessment of distress and interference related to traumatic exposure.

Psychometric Properties

Sample Scores and Norms. Normative information is available for a variety of samples that vary across gender, age, and trauma type derived from 10 studies reported by King, King, Leskin, and Foy (1995). *PTSD severity* scores for male veterans have been found to range from 46.94 to 49.82; in the same samples the *LASC total* score ranged from 94.63 to 107.87 (Leskin & Foy, 1993; Pava, 1993). Mean *PTSD severity* scores for help seeking women with a history of childhood sexual assault ranged from 29.57 to 31.18; in the same samples the *total LASC* scores ranged from 56.83 to 64.62 (Lawrence, 1992; Rowan, Foy, Rodriguez, & Ryan, 1994; Ryan, 1992).

Reliability. A data set was formed to assess the psychometric characteristics of the LASC by combining data from 10 studies that had used the measure with clinical samples (King et al., 1995). The samples were derived from a diverse set of populations including Vietnam veterans, battered women, adult survivors of childhood sexual abuse, maritally distressed women, psychiatric outpatients, and high-risk adolescents. The total data set included 874 respondents. Coefficient alpha was .94 for the 17 items specifically assessing PTSD symptoms and .95 for the total score. Test-retest reliability over 2 weeks was available for a sample of 19 Vietnam veterans. The 17-item scores yielded a coefficient of .94 and the total scores yielded a coefficient of .90.

Validity. A factor analysis on the combined data set yielded three factors accounting for a total of 40.8% of the variance in the scale. Factor 1 was represented primarily by items that assessed the specific symptoms of PTSD. Factor 2 included items that tapped into physical manifestations of stress (e.g., severe headaches, abdominal distress, dizziness). Factor 3 included items reflecting issues related to interpersonal functioning (e.g., marked selfconsciousness, inability to make and keep same-sex friends).

The LASC PTSD scores have been shown to be moderately but significantly related to measures of combat exposure (rs ranging from .30 to .51; Foy, Sipprelle, Rueger, & Carroll, 1984; Lund, Foy, Sipprelle, & Strachan, 1984; Resnick, Foy, Donahoe, & Miller, 1989) and other self-report measures of PTSD symptomatology (rs ranging from .38 to .48; Astin, Lawrence, & Foy, 1993). Diagnoses based on LASC scores corresponded to diagnosis based on a structured clinical interview with a sensitivity rate of 70% and a specificity rate of 80% (Housekamp & Foy, 1991).

King et al. (1995) also examined the ability of the LASC to predict PTSD diagnosis derived from a structured clinical interview. Although the 17 items were associated with a PTSD diagnosis (a score of 34 on the LASC is associated with approximately a 75% probability of having PTSD), the additional 26 items did not add to the predictive power of the measure. Thus, although these items may be clinically descriptive, their predictive validity is unproven.

Alternate Forms

An adolescent version of the LASC is available (Foy, Wood, King, King, & Resnick, 1997).

Source

The LASC is reprinted in Appendix B. The primary author of the scale is David W. Foy, Ph.D., Graduate School of Education and Psychology, Pepperdine University Plaza, 400 Corporate Pointe, Culver City, CA 90230, USA; (tel) 310-568-5739; (fax) 310-568-5755. More information can also be obtained by contacting Lynda A. King, National Center for PTSD (116B-3), VA Boston Healthcare System, 150 South Huntington Avenue, Boston, MA, 02130, USA; (tel) 617-232-9500, ext. 4938; (fax) 617-566-8508; (e-mail) lking@world. std.com.

MINNESOTA MULTIPHASIC PERSONALITY INVENTORY PTSD SCALE (MMPI-PTSD)

Original Citation

Keane, T. M., Malloy, P. F., & Fairbank, J. A. (1984). Empirical development of an MMPI subscale for the assessment of combat-related posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 52, 888–891.

Purpose

To detect symptoms of posttraumatic stress disorder.

Description

One of the greatest advantages of the MMPI-PTSD scale is that it is widely available to clinicians who regularly administer the MMPI-2 as part of their practice. The original MMPI-PTSD was a 49-item measure derived from the MMPI. When the MMPI was renamed and revised, the MMPI-PTSD scale also underwent some changes including the deletion of three items, the rewording of one item, and a change in item order (thus the current version has 46 items). The items are answered in a true-false format. Although the scale is typically administered as part of the full MMPI-2, it can be useful as a stand-alone scale. The embedded and stand-alone versions have been shown to be highly correlated (r = .90; Herman, Weathers, Litz, & Keane, 1996).

Administration and Scoring

It takes 15 minutes to administer the stand-alone version of the MMPI-PTSD scale. A total score is derived by summing the positive answers to the items. The MMPI-2 PTSD is also

MEASURES FOR ACUTE STRESS DISORDER AND PTSD 273

discussed in the MMPI-2 manual and can be scored by the hand or computer method. A cutoff score of 30 was originally suggested for detecting PTSD among veterans; later studies with the MMPI-2 version suggested a cutoff between 24 and 28. The civilian cutoff is in the range of 15 to 19. Scores greater than 38 or 40 may indicate fabrication of symptoms. Additional information about scoring and interpreting the MMPI-PTSD scale in the context of the MMPI-2 is available in the manual.

Psychometric Properties

Sample Scores and Norms. On the original MMPI-PTSD scale, veterans with PTSD have a mean score between 26 and 37 (Keane et al., 1984; Koretzky & Peck, 1990), whereas psychiatric controls have a mean of 20 (Keane et al., 1984) and non-PTSD patients a mean of 12.30 (Koretzky & Peck, 1990). Veterans with PTSD revealed a mean of 31.5 on the revised stand-alone scale whereas those without PTSD had a mean of 15.5 (Herman et al., 1996). Mean scores for the revised, embedded MMPI-PTSD scale have been shown to range from 30.6 to 36.2 for veterans with PTSD (Herman et al., 1996; Litz et al., 1991), and to be 15.5 for veterans with PTSD (Herman et al., 1996), 22.9 for psychiatric controls, 18.3 for substance abusers, and 5.2 for a comparison group (Litz et al., 1991).

Reliability. The internal consistency of the embedded and stand-alone versions of the MMPI-2 PTSD scale has been shown to be excellent in a veteran sample (α s range from .95 to .96; Herman et al., 1996).

The test-retest reliability of the stand-alone version of the MMPI-2 PTSD scale over 2 to 3 days was also excellent in a veteran sample (r = .95; Herman et al., 1996).

Validity. Veterans with PTSD have been shown to score significantly higher than psychiatric controls (Keane et al., 1984) and veterans without PTSD (Herman et al., 1996; Scotti, Sturges, & Lyons, 1996). Further, the MMPI-PTSD scale can differentiate veterans with comorbid PTSD and substance abuse from those with substance abuse alone (Kenderdine, Phillips, & Scurfield, 1992).

A score of 30 correctly classified 82% of veterans with and without PTSD (Keane et al., 1984). A score of 19 correctly classified 88% of patients in a psychotherapy clinic (Koretzky & Peck, 1990).

With regard to convergent validity, the original MMPI-PTSD scale was correlated significantly with other self-report measures of PTSD in a sample of veterans (rs ranged from .79 to .88; Watson et al., 1994) although another study found the associations to be lower (rs ranged from .21 to .71; McFall, Smith, Mackay, & Tarver, 1990). In a sample of battered women, Dutton, Perrin, Chrestman, Halle, and Burghardt (1991) found the MMPI-PTSD scale to be moderately but significantly correlated with other self-report measures of PTSD (rs ranged from .33 to .65). Stronger convergent validity with other measures of PTSD was demonstrated in a mixed trauma sample (Neal, Busuttil, Rollins, et al., 1994; rs ranged from .79 to .85); however, in this sample, the scale was also highly correlated with a measure of general distress (r = .92). Further, the MMPI-PTSD scale was significantly correlated with a number of MMPI scales and subscales in a sample of inpatient alcoholics, again suggesting that the measure may assess general psychological distress (Moody & Kish, 1989).

The embedded and stand-alone versions of the MMPI-2 PTSD scale have been shown to be moderately but significantly associated with a measure of combat exposure (rs ranged from .32 to .37), other self-report measures of PTSD (rs ranged from .65 to .85), and a structured clinical interview for PTSD (rs ranged from .77 to .80; Herman et al., 1996).

274 CHAPTER 19

Cutoffs of 26 and 28 for the embedded MMPI-2 PTSD scale have been shown to correctly classify 76% of veterans with PTSD (Herman et al., 1996; Munley, Bains, Bloem, & Busby, 1995), whereas a cutoff of 24 on the stand-alone version correctly classified 80% of veterans with PTSD (Herman et al., 1996).

Treatment sensitivity for the MMPI-PTSD has been demonstrated in a number of studies (e.g., Brom, Kleber, & Defares, 1989; Thompson et al., 1995).

Source

Additional information about the MMPI-PTSD scale can be obtained by contacting Terence M. Keane, Ph.D., National Center for PTSD (116B-2), VA Boston Healthcare System, 150 South Huntington Avenue, Boston, MA 02130, USA; (tel) 617-232-9500, ext. 4143; (e-mail) terry.keane@med.va.gov.

MISSISSIPPI SCALE FOR PTSD

Original Citations

Keane, T. M., Caddell, J. M., & Taylor, K. L. (1988). Mississippi Scale for Combat-Related Posttraumatic Stress Disorder: Three studies in reliability and validity. *Journal of Consulting and Clinical Psychology*, 56, 85–90.

Purpose

To measure self-reported symptoms of PTSD.

Description

There are several versions of the Mississippi Scale for PTSD. Among the most widely used include the Mississippi Scale for Combat-Related PTSD and the Civilian Mississippi Scale. The original combat scale consists of 35 items that tap into the presence of symptoms reflecting the three main DSM-IV criteria for PTSD: reexperiencing (criterion B), avoidance and numbing (criterion C), and hyperarousal (criterion D) and associated features (e.g., depression, substance abuse). Items are rated on a five-point scale with anchors that vary depending on the item but include phrases such as "not at all true" to "almost always true." Respondents are asked to rate symptoms over the time period occurring "since the event." The original version of the civilian scale used in the civilian/nonveteran component of the National Vietnam Veterans Readjustment Study (NVVRS) also had 35 items. Eleven of the items were rephrased slightly, changing reference to military service to a more general reference to the past. In both the combat and civilian versions, 4 items were added to make the scale consistent with DSM-IV criteria. These items assess symptoms of reexperiencing, psychogenic amnesia, hypervigilance, and increased arousal when confronted with reminders of the event. However, these items have not been found to increase the discriminative validity of the measure, so they are commonly omitted.

ADMINISTRATION AND SCORING

The full Mississippi Scale takes 10 to 15 minutes to administer. After reversing the positively worded items, a total score is derived by summing all of the items. A cutoff score of 107 was originally established for the combat version, although later studies suggested that a cutoff of 121 allows for better differentiation between veterans with and without PTSD. For the version reprinted in Appendix B, there are 9 positively worded items that should be reversed (items 6, 11, 17, 19, 22, 24, 27, 30, 34).

Psychometric Properties

Sample Scores and Norms. Among a large sample of treatment seeking veterans, the mean score for the 35-item scale was 104.5 (SD = 26.2; Keane et al., 1988). Means on the civilian measure for undergraduates ranged from 73.5 to 74.4 on the 35-item scale, and 81.8 to 82.9 on the 39-item scale (Lauterbach, Vrana, King, & King, 1997). The mean score obtained by civilians on the NVVRS was 64.3 (SD = 13.2) for the 35-item scale (Vreven, Gudanowski, King, & King, 1995).

Reliability. In a large sample of veterans seeking treatment, the 35-item Mississippi Scale for Combat-Related PTSD was shown to have excellent internal consistency ($\alpha = .94$; Keane et al., 1988). Test-retest reliability over 1 week in a smaller sample of veterans was .97 (Keane et al., 1988).

Internal consistency for the civilian version has also been demonstrated to be very good for both the 35- and 39-item scales (α s from .86 to .89; Lauterbach et al., 1997; Vreven et al., 1995).

Validity. Several factor analyses have been conducted on the 35-item Mississippi Scale for Combat-Related PTSD (e.g., Keane et al., 1988; McFall, Smith, Mackay, & Tarver, 1990). King and King (1994) conducted exploratory and higher order confirmatory factor analysis on the Mississippi Scale for Combat-Related PTSD using data from over 2200 veterans who participated in the NVVRS. The results suggest that the latent structure of this scale is best represented as an overarching single PTSD factor with four subsidiary dimensions: (1) reexperiencing and situational avoidance, (2) withdrawal and numbing, (3) arousal and lack of control, and (4) self-persecution.

Factor analyses on the civilian version have yielded mixed results from study to study and for the 35- versus the 39-item scale (Lauterbach et al., 1997; Vreven et al., 1995).

With regard to convergent validity, scores on the Mississippi Scale for Combat-Related PTSD are significantly associated with combat exposure (rs range from .25 to .44; Keane et al., 1988; McFall, Smith, Mackay, & Tarver, 1990; McFall, Smith, Roszell, et al., 1990) and other self-report measures of PTSD (rs range from .44 to .88; McFall, Smith, Mackay, & Tarver, 1990; McFall, Smith, Roszell, et al., 1990; Watson et al., 1994).

The convergent validity of the civilian measure has also been examined. Individuals with symptoms of PTSD score significantly higher than those without any PTSD symptoms, and some relationship has been established between exposure to stressful events and score on the Civilian Mississippi Scale (Lauterbach et al., 1997; Vrenen et al., 1995). Also, civilian scores have been associated with sexual abuse-related posttraumatic symptomatology (Gold & Cardeña, 1998). Further, the scores have been significantly but moderately associated with other measures of PTSD (rs range from .34 to .52; Lauterbach et al., 1997). However, the

civilian version has also been found to be strongly associated with general distress (r = .63; Vrenen et al., 1995), depression (r = .71), and anxiety (r = .70; Lauterbach et al., 1997). Overall these findings suggest that the civilian version may be more of a general measure of psychopathology than a specific measure of PTSD (Lauterbach et al., 1997; Vrenen et al., 1995).

Compared with a diagnosis derived by structured interview, the diagnostic accuracy of the Mississippi Scale for Combat-Related PTSD with a cutoff score of 107 was 90%, with a sensitivity of 93% and a specificity of 89% (Keane et al., 1988). McFall, Smith, Mackay, and Tarver (1990) found similar diagnostic efficiency with a lower cutoff score. However, Dalton, Tom, Rosenblum, Garte, and Aubuchon (1989) reported that 77% of nonveterans were able to feign a score on the scale exceeding the 107 cutoff. Lyons, Caddell, Pittman, Rawls, and Perrin (1994) also found that the scale was vulnerable to faking and suggested a cutoff of 121. However, although the sensitivity of this cutoff is good (.95), its specificity is relatively low (.45).

The diagnostic accuracy of the civilian version of the scale was examined in a large sample of individuals recruited from an emergency room (Shalev, Freedman, Peri, Brandes, & Sahar, 1997). With a cutoff of 75, the sensitivity of the scale was .87, and the specificity was .51. No cutoff score was found that could optimize both sensitivity and specificity.

Alternative Forms

The Mississippi Scale for Combat-Related PTSD has been translated into Hebrew and Spanish. A short version of the combat-related version, comprised of 11 items, has been shown to have good internal consistency, high sensitivity and specificity against a cutoff derived from the full scale, and good treatment sensitivity (Fontana & Rosenheck, 1994). Norris and Perilla (1996) developed a 30-item Revised Version of the Civilian Mississippi Scale that has demonstrated internal consistency for both an English and a Spanish translation.

Source

The Mississippi Scale for Combat-Related PTSD can be obtained by contacting Terence M. Keane, Ph.D., National Center for PTSD (116B-2), VA Boston Healthcare System, 150 South Huntington Avenue, Boston, MA 02130, USA; (tel) 617-232-9500, ext. 4143; (e-mail) terry.keane@med.va.gov. The 35-item Civilian Mississippi Scale is reprinted in Appendix B.

PENN INVENTORY FOR PTSD (PENN INVENTORY)

Original Citation

Hammarberg, M. (1992). Penn Inventory for Posttraumatic Stress Disorder: Psychometric properties. *Psychological Assessment*, 4, 67-76.

Purpose

To measure severity of PTSD.

Description

の方法を行いていたので

The Penn Inventory is a 26-item self-report measure of the severity of PTSD. Each item comprises four sentences modeled after the Beck Depression Inventory. The meanings of the series of sentences measure the presence or absence of PTSD symptoms over the past week in addition to their degree, frequency, or intensity. The respondent chooses the statement that best describes their experience. Each sentence is rated from 0 to 3 with higher scores representing more symptomatology. Items are not keyed to a specific traumatic event.

Administration and Scoring

The Penn Inventory can be administered in 10 to 15 minutes. A score is derived by summing all of the circled values. A cutoff score of 35 can be used to determine the likely presence of PTSD.

Psychometric Properties

Sample Scores and Norms. The mean scores for clinical samples of Vietnam veterans range from 51.1 (SD = 12.3) to 54.7 (SD = 8.7; Hammarberg, 1992). Veterans without PTSD revealed a mean score of 15.6 (SD = 9.1) and a nonveteran community sample had a mean of 15.3 (SD = 8.4; Hammarberg, 1992).

Reliability. The Penn Inventory has been demonstrated to have very good to excellent internal consistency (α ranges from .78 to .94) across a variety of clinical and community samples (Hammarberg, 1992). Additionally, test-retest over an average of 5.2 days ranged from .87 to .93 (Hammarberg, 1992).

Validity. Individuals diagnosed with PTSD on the basis of self-report measures and a clinical interview scored significantly higher than those without PTSD (either confirmed by diagnostic interview or assumed based on nonveteran or community status; Hammarberg, 1992).

The overall diagnostic efficiency of the measure against another self-report measure of PTSD among veterans was 94% (Hammarberg, 1992). In a sample of disaster victims, the hit rate was 95% (Hammarberg, 1992).

Scores on the Penn Inventory have been shown to be moderately but significantly associated with exposure to combat (r = .39; Hammarberg, 1992). Further, among a group of veterans diagnosed with PTSD, the Penn Inventory was demonstrated to be associated with measures of anxiety (rs ranged from .74 to .82) and depression (r = .52). The Penn Inventory was also significantly associated with additional measures of PTSD (rs from .72 to .85).

Treatment sensitivity was demonstrated in a sample of Vietnam veterans treated for PTSD (Hammarberg & Silver, 1994). Patients changed on average from a score of 55 (SD = 9.2) to a score of 45.96 (SD = 16.0) over a period of 12 weeks, whereas untreated veterans with PTSD and non-PTSD Vietnam era veterans and nonveterans showed no significant symptom change.

Source

The Penn Inventory may be obtained from Melvyn Hammarberg, Ph.D., Department of Anthropology, 325 Museum, University of Pennsylvania, Philadelphia, PA 19104-6398, USA; (tel) 215-898-0981; (fax) 215-898-7462; (e-mail) mhammarb@ecat.sas.upenn.edu. The cost of the measure is \$35.00 US.

POSTTRAUMATIC COGNITIONS INVENTORY (PTCI)

Original Citation

Foa, E. B., Ehlers, A., Clark, D. M., Tolin, D. F., & Orsillo, S. M. (1999). The Posttraumatic Cognitions Inventory (PTCI): Development and validation. *Psychological Assessment*, *11*, 303-314.

Purpose

To measure trauma-related thoughts and beliefs.

Description

The PTCI is a 36-item self-report questionnaire that taps into three constructs related to trauma-related thoughts and beliefs: *negative cognitions about self, negative cognitions about the world*, and *self-blame*. Each item presents a statement and is followed by a seven-point response scale representing degree of agreement ranging from 1 (totally disagree) to 7 (totally agree). Items are worded so that higher ratings reflect greater endorsement of pathological cognitions.

Administration and Scoring

The PTCI can be administered in 10 minutes. Scoring consists of summing the items that make up each subscale and dividing the sum by the number of items comprising the subscale. *Negative cognitions about self* is derived from items 2–6, 9, 12, 14, 16, 17, 20, 21, 24–26, 28–30, 33, 35, and 36; *negative cognitions about the world* is derived from items 7, 8, 10, 11, 18, 23, and 27; *self-blame* is derived from items 1, 15, 19, 22, and 31. Items 13, 32, and 34 are experimental and are not included in the subscales. The total score is derived by taking the sum of the items that comprise the three subscales.

Psychometric Properties

The psychometric properties are derived from a sample of 601 volunteers, 392 of whom had experienced a traumatic event, and 170 of whom had reported at least moderate PTSD symptomatology on a self-report measure (Foa et al., 1999). Participants were recruited from clinical, community, and undergraduate settings.

MEASURES FOR ACUTE STRESS DISORDER AND PTSD 279

Sample Scores and Norms. The median scores for a sample of individuals with PTSD of at least moderate severity were 133 (SD = 44.17) for total score, 3.60 (SD = 1.48) for negative cognitions about self, 5.00 (SD = 1.25) for negative cognitions about the world, and 3.20 (SD = 1.74) for self-blame. Median scores derived from a nontraumatized group were 45.50 (SD = 34.76) for total score, 1.08 (SD = 0.76) for negative cognitions about self, 2.07 (SD = 1.43) for negative cognitions about the world, and 1.00 (SD = 1.45) for self-blame.

Reliability. Cronbach's alphas for the three PTCI scales and total scales are good to very good (total score, $\alpha = .97$; negative cognitions about self, $\alpha = .97$; negative cognitions about the world, $\alpha = .88$; self-blame, $\alpha = .86$). One-week test-retest reliability on a subsample of the respondents was .74 for the total score and ranged from .75 to .89 for the scales. Three-week test-retest reliability in another subsample was .85 for the total score and ranged from .80 to .86 for the scales.

Validity. Factor analysis confirmed the existence of the three-factor structure. The first factor explained 48.5% of the variance, the second factor accounted for an additional 4%, and the third factor accounted for an additional 3.4%. The stability of the structure was validated across three samples.

PTCI scores were found to correlate with PTSD severity (r = .79), depression (r = .75), and anxiety (r = .75). The scales of the PTCI were significantly associated with similar scales assessing trauma-related cognitions. Traumatized individuals with PTSD scored significantly higher than traumatized individuals without PTSD and nontraumatized individuals on all of the PTCI scales. Further, the PTCI compared favorably with other measures of trauma-related cognitions for predicting PTSD. Scores on the PTCI scales classified 86% of the traumatized individuals correctly into those with and without PTSD with a sensitivity of .78 and a specificity of .93.

Source

The PTCI is reprinted in the original citation and in Appendix B. Additional information about the measure can be obtained by contacting Edna B. Foa, Ph.D., Center for the Study and Treatment of Anxiety, University of Pennsylvania, School of Medicine, 3535 Market Street, 6th floor, Philadelphia, PA 19104, USA; (tel) 215-746-3327; (fax) 215-746-3311; (e-mail) foa@mail.med.upenn.edu.

POSTTRAUMATIC DIAGNOSTIC SCALE (PDS)

Original Citation

Foa, E. B., Cashman, L. A., Jaycox, L., & Perry, K. (1997). The validation of a self-report measure of posttraumatic stress disorder: The Posttraumatic Diagnostic Scale. *Psychological Assessment*, 4, 445–451.

Purpose

To assess the DSM-IV diagnostic criteria and symptom severity of PTSD.

280 CHAPTER 19

Description

The PDS is a revised varsion of an earlier self-report scale entitled the PTSD Symptom Scale (Foa, Riggs, Dancu, & Rothbaum, 1993). The PDS consists of 49 items arranged into four sections. Part I includes a checklist of 12 traumatic events one could experience or witness. In Part 2, the event causing the most distress in the past month is chosen, described in more detail and referred to in subsequent questions. Criterion A is established via four questions that assess physical threat and feelings of helplessness related to the event. Part 3 includes 17 items corresponding to PTSD criteria B, C, and D that assess the frequency of each symptom in the past month on a four-point scale. Part 4 assesses criterion F with 9 items that determine impairment in major life areas (e.g., work, leisure) using a yes/no format.

Administration and Scoring

The PDS can be administered in 10 to 15 minutes, It may be scored by hand or by computer program. A number of scoring indices can be derived including PDS diagnosis, symptom severity score, number of symptoms endorsed, symptom severity rating, and level of impairment in functioning. Following DSM-TV enteria, the diagnosis of PTSD requires the presence of physical injury or perception of life threat; a sense of helplessness or terror during the event; endorsement of at least one reexpariencing (criterion B) symptom, three avoidance (criterion C) symptoms, and two arousal (criterion D) symptoms; duration of at least 1 month; and impairment in at least one area of life functioning. An index of PTSD severity is obtained by summing the 17 symptom items.

Psychometric Properties

The psychometric properties are derived from a sample of 248 volunteers recruited from several PTSD meatment centers as well as from non-treatment seeking populations who may be at high risk for trauma (e.g., staff at police stations, individuals at women's shelters; Foa et al., 1997).

Sample Scores and Norms. The mean scores for a sample of 128 individuals with PTSD were 33.59 (SD = 9.96) for total symptom severity, 8.95 (SD = 3.68) for reexperiencing, 13.63 (SD = 4.76) for avoidance, and 11.02 (SD = 3.53) for arousal. The non-PTSD group (N = 120) obtained a mean score of 12.54 (SD = 10.54) on the total scale, 3.64 (SD = 3.18) on the reexperiencing scale, 4.54 (SD = 4.83) on the avoidance scale, and 4.36 (SD = 3.97) on the arousal scale.

Reliability. The PDS has been shown to have excellent internal consistency overall (a = .92) and very good internal consistency for the symptom subscales (as ranging from .78 to .84). Additionally, repeated administration over 2 to 3 weeks yielded an 87% agreement rate (kappa = .74) between diagnoses and adequate stability in symptom severity (all rs = .77 to .85).

Validity. Satisfactory agreement was found between the diagnoses derived from the PDS and those obtained from a structured clinical interview (kappa of .65, 82% agreement). Sensitivity of the PDS was .89 and specificity was .75. Scores reflecting symptom severity on

MEASURES FOR ACUTE STRESS DISORDER AND PTSD | 281

the PDS correlated with another measure of PTSD (r = .78), a measure of anxiety (rs range from .73 to .74), and a measure of depression (r = .79). These correlations raise the issue of whether the PDS is a specific measure of PTSD or a more general measure of psychological distress. However, given the high comorbidity of PTSD with anxiety and mood disorders, and the symptom overlap between disorders, this pattern of findings is not surprising.

Alternative Forms

The PTSD Symptom Scale Interview (PSS-I) is an interview version of the PTSD Symptom Scale Self-Report Scale that was the predecessor of the PDS. Psychometric properties of this scale are available in Foa et al. (1993).

Source

The PDS is available from National Computer Systems, P.O. Box 1416, Minneapolis, MN 55440, USA; (tel) 800-627-7271; (webpage) www.ncs.com. The PDS on-line version requires the purchase of Microtest Q Assessment Systems Software with an annual licensing fee of \$89.00 US. Each assessment profile costs \$4.25 US for the first 50 reports. The penciland-paper starter kit (including 1 manual, 10 answer sheets, and 1 scoring sheet) is \$44.00 US. The reorder kit (50 answer sheets, 50 work sheets, and 1 scoring sheet) is \$117.00 US. Additional information about the measure can be obtained by contacting Edna B. Foa, Ph.D., Center for the Study and Treatment of Anxiety, University of Pennsylvania, School of Medicine, 3535 Market Street, 6th floor, Philadelphia, PA 19104, USA; (tel) 215-746-3327; (fax) 215-746-3311; (e-mail) foa@mail.med.upenn.edu.

PTSD CHECKLIST (PCL)

Original Citations

Weathers, F. W., Litz, B. T., Herman, D. S., Huska, J. A., & Keane, T. M. (1993, October). The PTSD checklist: Reliability, validity and diagnostic utility. Paper presented at the Annual Meeting of the International Society for Traumatic Stress Studies, San Antonio, TX.

Blanchard, E. B., Jones-Alexander, J., Buckley, T. C., & Forneris, C. A. (1996). Psychometric properties of the PTSD Checklist (PCL). *Behaviour Research and Therapy*, 34, 669-673.

Purpose

To assess PTSD symptom severity.

Description

The PCL is a 17-item inventory that assesses the specific symptoms of PTSD. The respondent is asked to rate how much the problem described in each statement has bothered

him or her over the past month on a five-point scale ranging from 1 (not at all) to 5 (extremely). The authors have also suggested that the time frame (e.g., last week) can be changed to accommodate the goals of the assessment.

Administration and Scoring

The PCL takes 5 to 10 minutes to administer. A total score is an indicator of PTSD symptom severity. Cutoff scores of 50 for military samples and 44 for nonmilitary samples have been proposed (see below). Although the authors originally suggested that a PTSD diagnosis could be derived by considering a score of 3 or higher as reflecting the presence of a particular symptom, and by following the DSM-IV diagnostic rules to determine the appropriate number and pattern of symptoms, Blanchard et al. (1996) caution against this approach (see below).

Psychometric Properties

Sample Scores and Norms. In a sample of combat veterans, those with a diagnosis of PTSD obtained a mean of 63.58 (SD = 14.14) and those without a diagnosis of PTSD obtained a mean of 34.40 (SD = 14.09; Weathers et al., 1993). Individuals with MVA-related PTSD scored 60.0 (SD = 9.4) and those without PTSD scored 26.6 (SD = 4.6). Sexual assault victims diagnosed with PTSD scored 55 (SD = 16.7) versus 22.8 (SD = 11.8) for the no-PTSD assault group (Blanchard et al., 1996). Additional sample means are available for mothers of cancer survivors (Manne, Du Hamel, Gallelli, Sorgen, & Redd, 1998) and for breast cancer survivors (Andrykowski, Cordova, Studts, & Miller, 1998).

Reliability. The PCL has been shown to have excellent internal consistency in Vietnam and Persian Gulf veterans, victims of motor vehicle accidents, and sexual assault survivors (rs ranging from .94 to .97; Blanchard et al., 1996; Weathers et al., 1993). Test-retest reliability over 2 to 3 days was .96 for the Vietnam veterans (Weathers et al., 1993).

Validity. A factor analysis on data derived from the Persian Gulf war veterans suggested that the items are best accounted for by a single factor (Weathers et al., 1993). In a Vietnam veteran sample, the PCL-M was significantly correlated with other measures of PTSD (rs range from .77 to .93) and a measure of combat exposure (r = .46; Weathers et al., 1993). Among Persian Gulf veterans, the PCL-M was significantly associated with another measure of PTSD (.85; Weathers et al., 1993).

Several studies have examined the diagnostic efficiency of the PCL. Weathers et al. (1993) found that at a cutoff of 50, the PCL-M predicted PTSD diagnosis derived from a structured clinical interview with a sensitivity of .82 and a specificity of .84. Blanchard et al. (1996) found the same cutoff yielded a sensitivity of .78 and a specificity of .86, and that a cutoff of 44 improved the sensitivity to .94 and specificity to .86 with an overall diagnostic efficiency of 90%. However, they also found variability in the most efficient cutoff score for each item (3 versus 4), thus they caution against the use of a score of 3 on a sufficient number of criterion B, C, and D symptoms to derive a diagnosis.

Additional diagnostic efficiency for cancer groups is found in Manne et al. (1998) and Andrykowski et al. (1998).

MEASURES FOR ACUTE STRESS DISORDER AND PTSD 283

Alternative Forms

There are several versions of the PCL: the PCL-Military, the PCL-S (which is tied to a specified stressor), and the PCL-C (which is not tied to a specific stressful event, but instead asks about "response to stressful life events"). A parent report on child symptoms is also available (PCL-PR).

Source

The PCL-C is reprinted in Appendix B. More information about the scale can be obtained from Frank Weathers, Ph.D., Department of Psychology, 226 Thach Hall, Auburn University, AL 36849, USA, (tel) 334-844-6495; (e-mail) weathfw@mail.auburn.edu.

PURDUE PTSD SCALE-REVISED (PPTS-R)

Original Citation

Lauterbach, D., & Vrana, S. (1996). Three studies on the reliability and validity of a self report measure of posttraumatic stress disorder. Assessment, 3, 17-25.

Purpose

To assess the frequency of each PTSD symptom.

Description

The PPTSD-R is a self-report measure comprised of 17 items corresponding to the symptoms found within PTSD criteria B, C, and D. Respondents rate the frequency of occurrence within the previous month of each item on a five-point scale ranging from 1 (not at all) to 5 (often).

Administration and Scoring

The PPTSD-R can be administered in 10 minutes. The scale can be scored to yield a dichotomous index reflecting the presence or absence of PTSD or to yield a continuous measure of severity. Continuous scores are obtained by summing the 17 items. The diagnosis of PTSD requires the endorsement of at least one *reexperiencing* (criterion B) symptom (items 1-4, 8), three *avoidance* (criterion C) symptoms (items 5-7, 9-12) and two *arousal* (criterion D) symptoms (items 13-17).

284 | CHAPIER 19

Psychometric Properties

The psychometric properties reported below are published in a multisample, multistudy paper (Lauterbach & Vrana, 1996).

Sample Scores and Norms. Mean scores for a sample of 440 undergraduate students are 31.5 (SD = 12.9) for the total score, 8.5 (SD = 4.1) for reexperiencing, 12.6 (SD = 5.5) for avoidance, and 10.4 (SD = 4.9) for arousal. Within a sample of 35 students receiving psychotherapy at a university-based counseling center the means were 38.7 (SD = 15.9) for the total score, 9.5 (SD = 4.6) for reexperiencing, 15.6 (SD = 6.7) for avoidance and 13.6 (SD = 6.3) for arousal.

Reliability. The PPTSD-R has excellent internal consistency overall ($\alpha = .91$) and very good internal consistency for the symptom subscales (α s ranging from .79 to .84). Test-retest reliability for 51 undergraduate students over 2 weeks reflected adequate stability in symptom severity for the total score (r = .72); however, stability was somewhat lower for the *avoidance* (r = .67), *arousal* (r = .71), and *reexperiencing* subscales (r = .48).

Validity. The PPTSD-R has been shown to be more strongly correlated with other measures of PTSD symptomatology (rs range from .50 to .66) than measures of anxiety (r = .37) and depression (r = .39). Further, students who experienced at least one traumatic event scored significantly higher on the PPTSD-R than those who did not report any traumatic events on the total score, *reexperiencing* and *arousal* subscales. Although the traumatized group scored higher on the *avoidance* scale as well, this difference did not reach conventional levels of significance.

Alternative Forms

There is a military and a civilian version of the measure.

Source

The PPTSD-R is reprinted in Appendix B. More information is available from Dean Lauterbach, Ph.D., 350 Sam Sibley Drive, Room 313, Bienvenu Hall, Department of Psychology, Northwestern State University, Natchitoches, LA 71497, USA; (tel) 318-357-5453; (fax) 318-357-6802.

SHORT SCREENING SCALE FOR PTSD

Original Citation

Breslau, N., Peterson, E. L., Kessler, R. C., & Schultz, L. R. (1999). Short screening scale for DSM-IV posttraumatic stress disorder. American Journal of Psychiatry, 156, 908-911.

Purpose

To screen for PTSD in persons exposed to a DSM-IV qualifying traumatic event.

Description

The Short Screening Scale for PTSD is a seven-item (yes/no format) clinicianadministered interview measure derived from the modified National Institute of Mental Health Diagnostic Interview Schedule and the World Health Organization Composite International Diagnostic Interview developed and used in the Detroit Area Survey of Trauma (Breslau, Kessler, & Peterson, 1998). The majority of PTSD symptom measures ask about symptoms in connection with only one event. Often, respondents have experienced multiple events, but they are asked to choose the worst or most distressing event to complete the measure. The Short Screening Scale for PTSD was developed to enable interviewers to quickly and efficiently assess PTSD in response to a number of traumas.

Administration and Scoring

The Short Screening Scale for PTSD can be administered in less than 3 minutes. Scoring consists of counting the number of positive answers to the seven items. A score of 4 or more seems to be the best cutoff for predicting PTSD diagnosis.

Psychometric Properties

Sample Scores and Norms. None are available.

Reliability. The reliability of the seven-item screening scale has not been directly examined. However, the reliability of the PTSD module from which it was derived has been assessed. A random sample of 32 baseline PTSD cases and 23 noncases was selected to be reassessed 12 to 18 months after the baseline interview (Breslau et al., 1998). There was agreement on 83% of the cases.

Validity. The predictive validity of the Short Screening Scale for PTSD relative to the full diagnostic interview was examined in a representative sample of 1830 men and women who were interviewed as part of the 1996 Detroit Area Survey of Trauma (Breslau et al., 1999). With 4 as a cutoff, the sensitivity was 80.3% and the specificity was 97.3%.

Although this measure appears to be a promising screening tool, particularly for researchers, it is important to note that the validity of the scale has not been examined in a study in which the seven items were administered as a freestanding scale.

Source

The Short Screening Scale for PTSD is reprinted in Appendix B. More information about the scale can be obtained by contacting Naomi Breslau, Ph.D., Henry Ford Health Systems, Psychiatry Service, 3A, Detroit, MI 48202-3450, USA; (tel) 313-876-2516; (fax) 313-874-6221; (e-mail) nbresla1@hfhs.org.

286 CHAPTER 19

STANFORD ACUTE STRESS REACTION QUESTIONNAIRE (SASRQ)

Original Citation

Cardeña, E., Koopman, C., Classen, C., Waelde, L. C., & Spiegel, D. (2000). Psychometric properties of the Stanford Acute Stress Reaction Questionnaire: A valid and reliable measure of acute stress. *Journal of Traumatic Stress*, 13, 719-734.

Purpose

To assess the psychological symptoms experienced in the aftermath of a traumatic event.

Description

The SASRQ is a 30-item self-report measure of ASD. The instructions allow the administrator to specify the time period during which the respondent's symptoms should be rated. The respondent is asked to describe the stressful event and rate how much disturbance it caused. Then, the respondent rates 30 items on a six-point scale ranging from 0 (not experienced) to 5 (very often experienced). Items tap into dissociation (10 items), reexperiencing (6 items), avoidance (6 items), anxiety and hyperarousal (6 items), and impairment in functioning (2 items). A final question asks the respondent how many days he or she experienced the worst symptoms of distress.

Administration and Scoring

The SASRQ can be administered in 15 minutes. It can be scored continuously by summing all of the items or dichotomously (ratings between 0 and 2 = 0, ratings between 3 and 5 = 1) for the presence of a symptom. To meet criterion B, a respondent must endorse three or more of the symptom criteria for dissociation: numbing (items 20, 28), reduction in awareness of surroundings (items 4, 24), derealization (items 3,18), depensionalization (items 10, 13), dissociative amnesia (items 16, 25). A respondent must endorse a symptom within each of the remaining criterion symptom clusters to obtain an ASD diagnosis: criterion C (items 6, 7, 15, 19, 23, 29), criterion D (items 5, 11, 14, 17, 22, 30), criterion E (items 1, 2, 8, 12, 21, 27), and criterion F (items 9, 26).

Psychometric Properties

The psychometric properties of an earlier version and of the final version of the SASRQ were evaluated together in a multisample, multistudy paper (Cardeña et al., 2000).

Sample Scores and Norms. In a sample of 43 adult emergency rescue workers the mean score was 26.37 (SD = 25.52). In contrast, within a group of 97 nonexposed rescue workers the mean was 4.91 (SD = 8.34).

Reliability. The internal consistency for the 30 symptom items of the SASRQ was found to be very good to excellent for the total score (rs range from .80 to .95) and acceptable to excellent for the subscales (rs range from .64 to .91).

Test-retest reliability over 3 to 4 weeks with a sample of students who had not experienced a severe stressor in the interim was .69.

Validity. In several samples, hypothesized group differences on the SASRQ were confirmed. For instance, participants exposed to a rescue operation scored significantly higher on the SASRQ than their nonexposed colleagues. Gulf War veterans with PTSD scored significantly higher than controls.

Convergent and discriminant validity have been demonstrated with the SASRQ. Scores on the dissociation and anxiety subscales of the SASRQ were significantly correlated with a measure of intrusion and avoidance (rs ranging from .55 to .75). The association of these subscales with a measure of schizophrenic symptoms was weaker (rs ranging from .22 to .47). In another sample, score on the SASRQ was significantly associated with a measure of peritraumatic dissociation (r = .72).

Some preliminary support exists for the predictive validity of the SASRQ. Among workers in a building where a mass shooting occurred, all participants who met criteria for ASD had significant PTSD symptoms 7 months later. Individuals diagnosed with ASD on the SASRQ had significantly greater odds of being considered a PTSD case, based on a self-report measure, at 6-month follow-up.

Source

The SASRQ is reprinted in Appendix B. Additional information can be obtained by contacting Etzel Cardeña, Ph.D., Department of Psychology and Anthropology, University of Texas-Pan American, 1201 West University Drive, Edinburg, TX 78639, USA; (tel) 956-381-3329, ext. 3323; (fax) 956-381-3333; (e-mail) ecardena@panam.edu.

STRESSFUL LIFE EVENTS SCREENING QUESTIONNAIRE (SLESQ)

Original Citation

Goodman, L. S., Corcoran, C., Turner, K., Yuan, N., & Green, B. L. (1998). Assessing traumatic event exposure: General issues and preliminary findings for the Stressful Life Events Screening Questionnaire. *Journal of Traumatic Stress*, 11, 521–542.

[≫] Purpose

To assess lifetime exposure to a variety of traumatic events.

Description

The SLESQ is a 15-item self-report measure of lifetime exposure to traumatic events. Respondents are asked whether they have experienced an event (e.g., life-threatening illness, physically forced sexual relations), and if they endorse the item, they are asked a series of questions about the nature of their exposure that vary from item to item (e.g., the age at the time of the event, injuries they may have received, the nature of the relationship with the perpetrator). In addition to answering the questions, the respondents are also asked to describe each event. Finally, respondents indicate whether multiple items refer to the same event, and if an event happened more than once, they are asked to describe the nature of the additional episodes of the event. Criterion A-2 is not assessed with this measure. Although the majority of psychometric data derive from a college sample, the SLESQ was developed for use with community samples, and the psychometric properties are currently being evaluated in samples of low-income and ethnic minority women (L. A. Goodman, personal communication, August 16, 2000).

Administration and Scoring

The SLESQ takes 10 minutes to administer for most respondents; those with multiple traumas may need up to 20 minutes. The authors suggest that some level of screening of the description of each endorsed event be conducted to ensure that the responses are appropriate and that they truly reflect criterion A events (rather than stressful, but not traumatic, events). Depending on the question of interest, users can score the scale in a number of ways (for instance, count the total number of endorsed events, or the number of events in a specific category such as interpersonal versus noninterpersonal events).

Psychometric Properties

Sample Means and Norms. In a large sample of college students, 72% of respondents reported exposure to at least one traumatic event (Goodman et al., 1998). The mean number of events endorsed by this sample was 1.83 (SD = 1.96). The prevalence of each event was reported by Goodman et al. (1998). In a sample of 2507 women recruited from six college campuses, 65% of the sample reported at least one event (ruling out other events that were coded as nontraumatic; Green et al., 2000).

Reliability. Two-week test-retest reliability in college students for the number of events endorsed at each time point was .89 (Goodman et al., 1998). Kappas for specific events ranged from .31 for attempted sexual assault to 1.0 for robbery or mugging. Four items fell below a kappa of .40: attempted rape, witness to a traumatic event, other serious injury or life-threatening situation such as military combat or living in a war zone, other frightening or horrifying event (Goodman et al., 1998).

Validity. Overall prevalence rates for many of the traumatic events reported in the Goodman et al. (1998) sample were consistent with rates found in other large samples (e.g., sexual assault rates consistent with Koss, Gidycz, & Wisniewski, 1987). Rates of traumatic events were consistent with expected gender differences; women were more likely to have experienced molestation and attempted sexual assault whereas men were more likely to have experienced adult physical assault, and other serious injury or life threat (Goodman et al., 1998).

In a large multisite study, scores on a measure assessing posttraumatic functioning were significantly higher for women with interpersonal traumas than those who had experienced noninterpersonal traumas or no reported events (Green et al., 2000). Women with multiple interpersonal trauma events had significantly higher means than women with a single event (Green et al., 2000).

The convergent validity of the SLESQ against a clinical interview was assessed in a sample of students (Goodman et al., 1998). The correlation between the total number of events endorsed was .77. Kappas for the specific events ranged from .26 for witnessed death/assault to .90 for life threatening illness. Six items fell below a kappa of .60 with increased reporting in the interview condition (Goodman et al., 1998).

To assess the ability of the SLESQ to detect criterion A events, Goodman et al. (1998) reviewed the responses to a randomly selected subsample of questionnaires and rated the descriptions against a conservative definition of criterion A. The first three authors made decisions about whether an event surpassed the threshold. Eighty-five percent of the events described in the SLESQ met the authors' severity threshold for a traumatic event (Goodman et al., 1998).

Source

The SLESQ is reprinted in the original article. More information about the measure can be obtained by contacting Lisa Goodman, Ph.D., Counseling Psychology Program, School of Education, Boston College, Chestnut Hill, MA 02467, USA; (tel) 617-552-1725; (e-mail) goodmalc@bc.edu.

STRUCTURED INTERVIEW FOR DISORDERS OF EXTREME STRESS (SIDES)

Original Citation

Pelcovitz, D., van der Kolk, B., Roth, S., Mandel, F., Kaplan, S., & Resick, P. (1997). Development of a criteria set and a structured interview for disorders of extreme stress (SIDES). *Journal of Traumatic Stress*, 10, 3–16.

Purpose

To measure alterations that may accompany exposure to extremely traumatic events.

Description

Several theorists and researchers have suggested that PTSD does not capture the full range of responses to traumatic events, particularly for traumatized children, rape victims, and battered women. Changes in affect regulation, self-identity, and interpersonal functioning seem to be common. The SIDES was developed to assess these proposed symptoms as part of the DSM-IV field trials. The SIDES consists of 48 items that tap into seven areas: (1) regulation of affect and impulses, (2) attention or consciousness, (3) self-perception, (4) perception of the perpetrator, (5) relations with others, (6) somatization, and (7) systems of meaning. There are also 27 subscales. Items have been described as being scored dichotomously

(Pelcovitz et al., 1997) and rated on a four-point scale ranging from "none or no problem with symptom" to "extremely problematic" (Zlotnick & Pearlstein, 1997).

Administration and Scoring

The original article developed criteria for determining the number of items in each subscale and scale necessary to meet a diagnostic cutoff (Pelcovitz et al., 1997).

Psychometric Properties

Reliability. In a sample of 520 community participants and treatment seekers who were part of the DSM-IV field trials, internal consistency for the scales ranged from .53 to .90 and was .96 for the total scale (Pelcovitz et al., 1997). Interrater reliability was established within a subsample of the community sample. The kappa coefficient rating for lifetime disorders was .81 (Pelcovitz et al., 1997).

Cronbach's alpha in a sample of survivors of childhood sexual assault was .90 for current (last 6 months) diagnosis and ranged from .42 to .84 for the individual scales (Zlotnick & Pearlstein, 1997).

Validity. The validity of the SIDES was examined in a sample of childhood sexual assault survivors. The *affect regulation* scale was correlated with a measure of borderline personality (r = .45), avoidance (r = .71), hypervigiliance (r = .50), impulsivity $(r^4 = .50)$, hostility (r = .55), and somatization (r = .51). Alterations in attention and consciousness was correlated with dissociation (r = .60) and avoidance (r = .57). Alterations in self-perception was correlated with avoidance (r = .56), disconnection (r = .51), and borderline personality (r = .46). Somatization was correlated with another measure of somatization (r = .68), avoidance (r = .63), and hypervigilance (r = .57). None of these scales were significantly correlated with a measure of narcissism (Zlotnick & Pearlstein, 1997).

Source

The SIDES can be obtained by contacting David Pelcovitz, Ph.D., Department of Psychiatry, North Shore University Hospital-Cornell University Medical College, Manhassett, NY, 11030, USA.

STRUCTURED INTERVIEW FOR PTSD (SIP)

Original Citations

Davidson, J., Smith, R., & Kudler, H. (1989). Validity and reliability of the DSM-III criteria for posttraumatic stress disorder: Experience with a structured interview. *Journal of Nervous and Mental Disease*, 177, 336–341.

Davidson, J. R. T., Malik, M. A., & Travers, J. (1997). Structured Interview for PTSD (SIP): Psychometric validation for DSM-IV criteria. *Depression and Anxiety*, 5, 127–129.

Purpose

To assess diagnostic status and symptom severity of PTSD.

Description

The SIP was originally developed to capture DSM-III symptoms or PTSD, but it has been revised and updated for DSM-IV. The SIP consists of 17 items representing the DSM-IV criteria for PTSD along with two measures of survivor and behavior guilt. Each item is rated on a five-point scale ranging from 0 (not at all) to 4 (extremely severe, daily or produces so much distress that patient cannot work or function socially).

Administration and Scoring

The SIP takes 10 to 30 minutes to administer depending on the complexity and severity of the individual's symptoms. The SIP can be scored using DSM-IV diagnostic criteria to yield a dichotomous score reflecting the presence or absence of the diagnosis, or the items can be summed to obtain a measure of symptom severity.

Psychometric Properties

Sample Scores and Norms. Among a sample of patients diagnosed with PTSD and enrolled in a clinical trial, the mean pretreatment SIP score was 36 (SD = 9.7; Davidson et al., 1997).

Reliability. Cronbach's alpha was .94 for a veteran sample (Davidson et al., 1989) and .80 for the clinical trial sample described above (Davidson et al., 1997). Interrater reliability on a subsample from the clinical trial ranged from .97 to .99 for total SIP score, for symptoms over the past 4 weeks and during the worst ever period (Davidson et al., 1997).

Test-retest reliability for a subsample of patients in the clinical trial who showed no clinical change on an independent measure of functioning between weeks 2 and 4 of the trial was .71 (Davidson et al., 1997). Test-retest reliability in a subsample of the veterans who demonstrated no change between weeks 4 and 8 of a clinical trial was .89 (Davidson et al., 1989).

Validity. A factor analysis conducted on an earlier version of the SIP revealed three factors representing arousal and intrusiveness (which accounted for the majority of the variance), guilt and avoidance, and problems with sleep, concentration, and numbing (Davidson et al., 1989). A factor analysis on the revised scale, conducted with a mostly female, chronic PTSD sample, yielded seven factors, with one strong factor accounting for most of the variance, and six weaker factors, some comprised of only a single item (Davidson et al., 1997).

The SIP has been shown to be significantly correlated with other measures of PTSD (rs range from .49 to .67, Davidson et al., 1989, 1997), but not with measures of combat exposure (r = .08) or intensity of combat (r = .27; Davidson et al., 1989). The SIP is also associated with interview measures of anxiety (rs range from .48 to .51) and depression (rs range from .42 to .57; Davidson et al., 1989, 1997). A small but significant relationship was found between

several measures of disability and the SIP (rs range from .25 to .33), but not social support (r = .14; Davidson et al., 1997).

The diagnostic sensitivity of the SIP as relative to the Structured Clinical Interview for DSM was 96% and the specificity was 80% (Davidson et al., 1989). At a score of 25, the SIP showed an efficiency of 94% relative to a structured interview diagnosis (Davidson et al., 1997).

The SIP showed treatment sensitivity in that those with PTSD following treatment in a clinical trial had higher scores than patients who no longer met diagnostic criteria (Davidson et al., 1997).

Alternative Forms

Carlier, Lamberts, Van Uchelen, and Gersons (1998) describe the development and validation of the Self-Rating Scale for PTSD, an abridged self-report version of the SIP. The scale is reprinted in the Carlier et al. (1998) article. The TOP-8, a scale derived from the SIP, includes the items that occurred most frequently and that demonstrated the most change in response to treatment (Davidson & Colket, 1997). The TOP-8 can be administered in 5 to 10 minutes and has been shown to have very good to excellent internal consistency, test-retest reliability, interrater reliability, convergent validity, and treatment sensitivity. More information about the TOP-8 can be obtained by contacting Jonathan Davidson, M.D. (see contact information below).

Source

The SIP can be obtained by contacting Jonathan R. T. Davidson, M.D., Department of Psychiatry, Duke University Medical Center, Box 3812, Durham, NC 27710, USA; (tel) 919-684-2880; (fax) 919-684-8866; (e-mail) tolme@acpub.duke.edu.

TRAUMA-RELATED GUILT INVENTORY (TRGI)

Original Citation

Kubany, E. S., Haynes, S. N., Aubueg, F. R., Manke, F. P., Brennan, J. M., & Stahura, C. (1996). Development and validation of the Trauma-Related Guilt Inventory. *Psychological Assessment*, 8, 428–444.

Purpose

To assess cognitive and emotional aspects of guilt associated with exposure to a traumatic event.

A. 10 Car

Description

The TRGI is a 32-item self-report questionnaire designed to measure guilt associated with the experience of a traumatic event. It includes three scales—global guilt, distress, and guilt cognitions—and three subscales (all part of guilt cognitions)—hindsight-bias/ responsibility, wrongdoing, and lack of justification.

Psychometric Properties

Psychometric data are all derived from a multistudy, multisample report by Kubany et al. (1996).

Sample Scores and Norms. Means for each of the three scales and subscales are available for the TRGI on a sample of 325 college students, 168 battered women, and 74 veterans (Kubany et al., 1996).

Reliability. Internal consistency in a sample of 100 women receiving services from a battered women's shelter was .86 to .90 for the scales and .67 to .82 for the subscales. Similar results were found for combat veterans.

In a sample of 32 students, 1-week test-retest reliability ranged from .73 to .86 for the scales and from .74 to .83 for the subscales. Similar results were found for veterans over an average of 8.4 days (SD = 6.01).

Validity. Three factor-analytic studies were conducted (on students with a history of traumatic events and battered women) to refine the TRGI and determine its factor structure. The final factor structure that best accounts for the data seems to be a four-factor solution that consists of a distress factor and three cognitive factors termed hindsight-bias/responsibility, wrongdoing, and lack of justification.

Convergent validity was demonstrated with a traumatized student sample in that the global guilt scale correlated with a measure of PTSD (r = .48) and depression (r = .60). The guilt cognitions subscale was correlated moderately with PTSD (r = .32) and depression (r = .32) and the distress scale was correlated .77 with PTSD and .59 with depression. Correlations for the subscales were smaller, and no relationship was found between lack of justification and depression or PTSD.

Convergent validity was also demonstrated for the veteran and battered women groups with scores on each of the TRGI measures correlated highly with other measures of guilt, PTSD, depression, self-esteem, shame, and social anxiety. Some discriminant validity was demonstrated for both the battered women and veteran groups in that the TRGI was only weakly associated with a measure assessing guilt over commonplace events.

Alternative Forms

The TRGI was validated and cross-validated with an ethnically diverse sample. A translation into Tagalog is available.

Source

The TRGI can be obtained by contacting Edward S. Kubany, Ph.D., National Center for PTSD, Department of Veterans Affairs, 1132 Bishop Street, Suite 307, Honolulu, HI 96813, USA; (tel) 808-566-1651; (fax) 808-566-1885; (e-mail) kubany@pixi.com. A published version of the TRGI is in development with Western Psychological Services, 12031 Wilshire Boulevard, Los Angeles, CA 90025, USA; (tel) 800-648-8857; (fax) 310-478-7838; (e-mail) custsvc@wpspublish.com; (website) www.wpspublish.com.

TRAUMA SYMPTOM INVENTORY (TSI)

Original Citation

Briere, J. (1995). Trauma Symptom Inventory professional manual. Odessa, FL: Psychological Assessment Resources.

Briere, J., Elliot, D. M., Harris, K., & Cotman, A. (1995). Trauma Symptom Inventory: Psychometrics and association with childhood and adult victimization in clinical samples. *Journal of Interpersonal Violence*, 10, 387–401.

Purpose

To evaluate acute and chronic posttraumatic symptomatology.

Description

The TSI is unique from many other measures of posttraumatic symptomatology in that it includes symptoms beyond those typically associated with PTSD and ASD such as other intraand interpersonal problems that often arise in individuals with a history of chronic psychological trauma. The TSI is a 100-item self-report scale that is comprised of 3 validity scales and 10 clinical scales. The validity scales assess underendorsement, overendorsement, and random or inconsistent response style. The clinical scales include *anxious arousal*, *depression*, *anger/ irritability*, *intrusive experiences*, *defensive avoidance*, *dissociation*, *sexual concerns*, *dysfunctional sexual behavior*, *impaired self-reference*, and *tension reduction behavior*. Respondents are asked to indicate how often each symptom has occurred within the past 6 months on a four-point scale ranging from 0 (never) to 3 (often).

- **S**

Administration and Scoring

The TSI can be administered in 20 minutes, except in the case of significantly clinically impaired patients. Raw scores are derived by summing the items that comprise each scale. Directions are included in the manual for the handling of responses that are left blank. Raw scores are converted to T scores using the appropriate profile for the respondent based on age and gender. Interpretation of the respondent's profile is described in the professional manual using each elevated scale and common two-point profiles.

No.

Psychometric Properties

Sample Scores and Norms. A normative sample is described in the professional manual including norms for male and female Navy recruits (Briere, 1995). Means and standard deviations are also available for males and females with and without a victimization history from a sample of 370 patients recruited from inpatient and outpatient treatment facilities (Briere et al., 1995).

Reliability. The TSI has been demonstrated to have very good to excellent internal consistency for the individual scales (α s ranging from .74 to .90) in a sample of 370 patients (Briere et al., 1995), and similar reliability was demonstrated in a university sample, the standardization sample, and a Navy recruit sample of 3569 (Briere, 1995). No test-retest reliability data are available.

Validity. Exploratory factor analysis in the standardization sample yielded two independent factors labeled "Generalized Trauma and Distress" and "Self-Dysfunction" (Briere, 1995). Similar results were obtained with the clinical sample (Briere et al., 1995). Confirmatory factor analysis supported a three-factor model theoretically developed by the author with the dimensions of posttraumatic stress, self-dysfunction, and dysphoric mood.

Individuals with a history of physical or sexual abuse scored significantly higher than those who did not report such a history on all 10 clinical scales of the TSI (Briere et al., 1995). In discriminant function analyses on the normative sample, the experience of adult and childhood interpersonal violence and disaster was associated with elevated TSI scores (Briere, 1995). Data bearing on the convergent and discriminant validity of the clinical scales are published in the manual, along with data supporting the incremental and criterion validity of the scale (Briere, 1995). For instance, in a subsample of the standardization sample (N = 449), an optimally weighted combination of TSI scales correctly predicted 92% of the true positive and 91% of the true negative cases of a self-report derived diagnosis of PTSD.

Alternative Forms

The TSC-40 is a similar scale that predates the TSI and is designed only for use in research (Briere & Runtz, 1989). The TSI-A is an 86-item alternate version of the TSI that contains very few sexual items (items from the *sexual concerns* or *dysfunctional sexual behavior* scales are dropped). There is a Trauma Symptom Checklist for Children that is somewhat comparable. The TSI has been translated into other languages for research purposes only.

Source

The TSI is available from Psychological Assessment Resources, Inc., P.O. Box 998, Odessa, FL 33556, USA; (tel) 800-331-TEST or 813-968-3003; (webpage) www.parinc.com. A TSI introductory kit that includes the manual, 10 item booklets, 25 hand-scorable answer sheets, and 25 each of the male and female profile forms is priced at \$89.00 US. A computerized scoring program of the TSI is available that provides raw scores, T scores, and a profile of the scales for a one time cost of \$199.00 US. Interpretation of the profile is not included. Information about the scale is also available from John Briere, Ph.D., Department of Psy-

chiatry, USC School of Medicine, 1934 Hospital Place, Los Angeles, CA 90033, USA; (tel) 323-226-5697; (fax) 323-226-5502; (e-mail) jbriere@hsc.usc.edu.

TRAUMATIC EVENTS QUESTIONNAIRE (TEQ)

Original Citation

Vrana, S., & Lauterbach, D. (1994). Prevalence of traumatic events and post-traumatic psychological symptoms in a non-clinical sample of college students. *Journal of Traumatic Stress*, 7, 289–302.

Purpose

To assess the frequency, type, and severity of trauma experienced.

Description

The TEQ is a self-report instrument that assesses experiences with 11 specific types of trauma selected from DSM-III-R and the relevant literature as potentially eliciting posttraumatic symptoms. Two residual categories are also included, in which respondents can record additional events that are not included in the list, and events that the individual feels are so traumatic that they cannot be discussed. For each event endorsed, respondents are asked the number of times the event occurred, their age at the time of the event, and on a seven-point scale ranging from 1 (not at all) to 7 (extremely), whether they were injured at the time, and how traumatic it is presently. Finally, respondents who endorsed more than one item are asked to indicate which event was the worst for them. Those who do not endorse any of the items in the scale are asked to describe the worst, stressful experience that they have encountered.

Administration and Scoring

The TEQ can be administered in 10 minutes. Several indices of traumatic exposure can be obtained including the total number of experiences (sum all "yes" responses to item A; continuous events such as sexual abuse are counted as one event) and severity of experiences (sum all responses to items C through F).

Þ

Psychometric Properties

The psychometric properties described below were derived from a sample of 440 college students (Vrana & Lauterbach, 1994).

Sample Scores and Norms. Eighty-four percent of the sample reported at least one traumatic event, and approximately one-third of the sample reported exposure to four or more individual events. Males experienced significantly more events than females.

L'ARGE

Reliability. No published data are available.

Validity. Respondents with at least one traumatic event reported significantly more depression, anxiety, and PTSD symptomatology than those who did not report exposure to any traumatic events. Further, the number of traumatic events was a significant predictor of depression, anxiety, and PTSD symptom severity.

Alternative Forms

There is a military and a civilian version of the TEQ.

Source

The TEQ (civilian version) is reprinted in Appendix B. More information is available from Dean Lauterbach, Ph.D., 350 Sam Sibley Drive, Room 313, Bienvenu Hall, Department of Psychology, Northwestern State University, Natchitoches, LA 71497, USA; (tel) 318-357-5453; (fax) 318-357-6802.

TRAUMATIC LIFE EVENTS QUESTIONNAIRE (TLEQ)

Original Citation

Kubany, E. S., Haynes, S. N., Leisen, M. B., Owens, J. A., Kaplan, A. S., Watson, S. B., & Burns, K. (2000). Development and preliminary validation of a brief broad-spectrum measure of trauma exposure: The Traumatic Life Events Questionnaire. *Psychological Assessment*, 12, 210–224.

Purpose

To assess exposure to a broad range of potentially traumatic events.

Description

The TLEQ is a self-report scale that assesses exposure to 21 types of potentially traumatic events. An open-ended question that assesses exposure to some other life threatening or highly disturbing events is also included. The events are described in behaviorally descriptive terms, and emotionally charged terms such as *rape* or *abuse* are avoided. For each event, respondents are asked to indicate the number of times the event occurred on a seven-point scale ranging from never to five times or more. Respondents are asked whether each event evoked fear, helplessness, or horror. Additional questions ask about physical injury and immediate emotional response to the event. Finally, in the case of exposure to multiple events, respondents are asked to indicate which event they perceive as the worst.

Administration and Scoring

The TLEQ can be administered in 10 to 15 minutes.

Psychometric Properties

Psychometric properties were evaluated in a multisample, multistudy report (Kubany et al., 2000).

Sample Scores and Norms. Reports of occurrences of traumatic events across five samples are presented by Kubany et al. (2000).

Reliability. The temporal stability of the TLEQ was evaluated in a sample of 42 battered women. Kappa coefficients assessing agreement over 2 weeks were above .40 for 20 of the 21 items and .60 or above for 12 items. The overall mean percentage of test-retest agreement was 86%. Correlations of frequency of occurrence reports for each event (with the exception of combat) averaged .77 and ranged from .50 to .93. The test-retest reliability of the occurrence or nonoccurrence of criterion A-2 (intense fear, helplessness, and horror) yielded kappa coefficients of .40 or higher for 19 of the 21 items and .60 or higher for 16 items. The overall percentage of agreement for criteria A-2 was 89%.

Validity. Seven experts in the area of PTSD rated relevance and representativeness of several aspects of the TLEQ (e.g., response format, individual items) for measuring PTSD as defined in DSM-IV. Overall item wording was rated very positively, as was adequacy of coverage for traumatic events.

Convergent validity was assessed in a sample of 62 undergraduate students. Agreement was evaluated between a self-report and interview version of a slightly modified version of the TLEQ. Kappa coefficients were .40 for 15 of the 16 items and above .60 for 13 items. The overall mean percentage of agreement was 92%. Convergent validity was also assessed for the two formats administered 1 week apart with similar, but slightly lower, agreement.

All but one Vietnam veteran with documented service records endorsed exposure to combat on the TLEQ. Among the battered women, 98% of those who met the cutoff for PTSD, indicated that they had experienced partner abuse with fear, helplessness, and horror at the time of the event on the TLEQ.

Discriminative validity was assessed in a sample of battered women. Women who met the cutoff for PTSD on a self-report measure reported significantly more types of traumatic events, more total events, and more events that evoked intense fear, helplessness, and horror on the TLEQ than women who did not meet the PTSD cutoff.

Ť

Alternative Forms

A computerized version is currently being validated in a grant-funded project.

Source

The TLEQ is available from Edward S. Kubany, Ph.D., National Center for PTSD, Department of Veterans Affairs, 1132 Bishop Street, Suite 307, Honolulu, HI 96813, USA; (tel)

MEASURES FOR ACUTE STRESS DISORDER AND PTSD 299

ine.

808-566-1651; (fax) 808-566-1885; (e-mail) kubany@pixi.com. A published version of the TLEQ is in development with Western Psychological Services, 12031 Wilshire Boulevard, Los Angeles, CA 90025, USA; (tel) 800-648-8857; (fax) 310-478-7838; (e-mail) custsvc@ wpspublish.com; (website) www.wpspublish.com.

BRIEF DESCRIPTIONS OF ADDITIONAL MEASURES

Evaluation of Life-time Stressors (ELS)

The ELS is a comprehensive questionnaire and interview protocol designed to assess exposure to potentially traumatic events across the life span. The ELS-Q is a 53-item screening questionnaire that addresses exposure to traumatic events in addition to behavioral correlates of a wide range of potentially traumatic events. For each event, respondents indicate "yes," "no," "I'm not sure," or "It happened to someone I knew." All nonnegative responses on the questionnaire are followed up on the ELS-I (interview). Initial psychometrics on the protocol are quite promising. The ELS-Q is not used as a stand-alone questionnaire. The ELS can be obtained from Karen E. Krinsley, Ph.D., PTSD 116B-2, VA Boston Healthcare System, 150 South Huntington Avenue, Boston, MA 02130, USA; (e-mail) krinsley.karen@boston.va.gov.

Harvard Trauma Questionnaire (HTQ)

The HTQ is a self-report scale designed to measure trauma and torture events and to assess symptoms of PTSD and current functioning among individuals affected by torture, trauma and war-related violence. It is available in over 30 languages including Cambodian, Vietnamese, Laotian, Bosnian, Croatian, Ethiopian, Spanish, and Japanese. Cultural validity, reliability, and the instrument's psychometric properties have been established in a number of culturally diverse settings. The measure and more information can be obtained from Richard F. Mollica, M.D., M.A.R., Harvard Program in Refugee Trauma, 8 Story Street, 3rd floor, Cambridge, MA 02138, USA; (tel) 617-496-5550; (fax) 617-496-5530; (e-mail) rmollica@ hprt.harvard.edu.

Life Stressor Checklist-Revised (LSCL-R)

The LSCL-R is a measure of lifetime exposure to stressful and traumatic events. It is still under development and the authors request that those who wish to use it consult with them first. The scale is reprinted in a paper by Wolfe and Kimerling (1997). For more information about the scale, contact Rachel Kimerling, Ph.D., Assistant Adjunct Professor, Department of Psychiatry, UCSF School of Medicine, San Francisco General Hospital, 1001 Potrero Ave., Suite 2100, San Francisco, CA 94110, USA; (tel) 415-206-6447; (fax) 415-206-3855; (e-mail) rachelk@itsa.ucsf.edu.

Modified PTSD Symptom Scale (MPSS-SR)

The MPSS-SR is a 17-item self-report measure designed to assess the frequency and severity of PTSD symptoms corresponding to DSM-III-R criteria. It was developed from the

Foa et al. (1993) PTSD Symptom Scale. Updates to the scale were made to be consistent with DSM-IV criteria, but this updated version is not published. The measure was validated against the Structured Clinical Interview for DSM in the DSM-IV field trials and some promising psychometric data are discussed in the original reference by Falsetti, Resnick, Resick, and Kilpatrick (1993). More information about the scale is available from Sherri Falsetti, Ph.D., National Crime Victims Center, Medical University of South Carolina, 171 Ashley Avenue, Charleston, SC 29425, USA; (tel) 843-792-2945; (fax) 843-792-3388; (e-mail) falsetsa@ musc.edu.

100

Peri-Traumatic Dissociative Experiences Questionnaire (PDEQ)

The PDEQ is a 10-item self-report measure of dissociation at the time of a traumatic event. A recently completed meta-analysis of PTSD predictors found that greater dissociation at the time of trauma, as assessed with the PDEQ, was the most robust predictor of current PTSD (Ozer, Best, Lipsey, & Weiss, manuscript under review). A rater version of the scale is also available. Both versions are available in Marmar, Weiss, and Metzler (1997). For more information about the scale, contact Charles R. Marmar, M.D., Langley Porter Institute, 401 Parnassus Avenue, Box F 0984, San Francisco, CA 94122, USA; (tel) 415-750-2126; (fax) 415-221-6347; (e-mail) marmar@itsa.ucsf.edu.

Personal Beliefs and Reactions Scale (PBRS)

The PBRS is a self-report scale that was developed to assess cognitive schemas that are often disrupted in the aftermath of sexual trauma. It consists of 40 items and eight scales: esteem, trust, self-blame, safety, intimacy, control, negative rape beliefs, and undoing. Preliminary psychometrics on the scale appear promising. The scale can be obtained by contacting Mindy B. Mechanic, Ph.D., Department of Psychology, University of Missouri-St. Louis, 8001 Natural Bridge Road, St. Louis, MO 63121, USA; (e-mail) mbmechanic@umsl.edu.

Posttraumatic Dissociation Scale (PTDS)

The PTDS is a 24-item self-report measure that assesses the frequency of occurrences of derealization, depersonalization, gaps in awareness, amnesia, and gaps in awareness accompanied by reexperiencing over the previous week. The measure is currently under development and has not yet been published. More information about the scale can be obtained from Eve Carlson, Ph.D., National Center for PTSD (352-117 MP), VA Palo Alto Health Care System, 795 Willow Road, Menlo Park, CA 94025, USA; (tel) 650-493-5000, ext. 24058; (fax) 650-617-2684; (e-mail) carlson@icon.palo-alto.med.va.gov.

PTSD Interview

The PTSD Interview is based on DSM-III-R criteria of PTSD. To date, it has been used primarily with veterans. A description of the measure and some psychometric data are available in a paper by Watson, Juba, Manifold, Kucala, and Anderson (1991). More information about the scale can be obtained from Dr. Charles Watson, VA Medical Center, Research Service, 4801 8th Street North, St. Cloud, MN 56303, USA.

PTSD Inventory

The PTSD Inventory is a 17-item self-report diagnostic questionnaire that assesses warrelated PTSD. The measure has been demonstrated to have very good validity relative to a structured interview. A revised version of the scale, consistent with DSM-IV criteria, is described and reprinted in Solomon et al. (1993). A Hebrew version of the scale is available from the author. More information about the scale can be obtained from Zahava Solomon, # Ph.D., Bob Shapell School of Social Work, Tel Aviv University, Ramat Aviv, Tel Aviv, Israel.

SCL-90-R PTSD Scale

A 28-item scale derived from the SCL-90-R has been developed and validated to detect crime-related PTSD (Saunders, Arata, & Kilpatrick, 1990). This scale was not designed to be administered independent of the SCL-90, instead it is interpreted in the context of the entire measure. Similarly, a 25-item war-zone-related PTSD scale with strong psychometric properties has been derived from the SCL-90-R (Weathers et al., 1996). The item numbers that make up these measures and the psychometric properties are available in the respective papers. More information about the crime-related PTSD scale is available from Benjamin Saunders, Ph.D., National Crime Victims Center, Medical University of South Carolina, 171 Ashley Avenue, Charleston, SC 29425, USA; (tel) 843-792-2945; (fax) 843-792-3388; (e-mail) saunders@ musc.edu. More information about the war-zone-related PTSD scale can be obtained from Frank Weathers, Ph.D., Department of Psychology, 226 Thach Hall, Auburn University, Auburn, AL 36849, USA; (tel) 334-844-6495; (e-mail) weathfw@mail.auburn.edu.

Self-Rating Inventory for Posttraumatic Stress Disorder (SIP)

The SIP is a 47-item Dutch language self-rating questionnaire that assesses PTSD symptoms and associated features. Psychometric data on the scale are found in Hovens, van der Ploeg, Klaarenbeek, Schreuder, and Rivero (1994b). More information is available from J. E. Hovens, Ph.D., Centre '45, National Center for the Treatment of WWII Victims, Hermanstraat 6, NL-2315 KS, Leiden, the Netherlands.

Stress Response Rating Scale (SRRS)

The SRRS is a clinician rating scale that assesses response to serious life-events. It consists of 40 items that reflect signs and symptoms of intrusion, avoidance, and distress. Symptoms are rated over the previous 7 days. More information about this measure and its psychometric properties are available in Weiss, Horowitz, and Wilner (1984). The scale is available from Daniel S. Weiss, Ph.D., Department of Psychiatry, University of California–San Francisco, Box F-0984, San Francisco, CA 94143, USA; (tel) 415-476-7557; (fax) 415-502-7296; (e-mail) dweiss@itsa.ucsf.edu.

Trauma Assessment for Adults (TAA)

There is a self-report and an interview version of the TAA, a brief screen for exposure to traumatic events. It is adapted from the Potential Stressful Events Interview (Falsetti, Resnick,

Kilpatrick, & Freedy, 1994) and the National Women's Study Event History and PTSD Module. More information about the scale can be obtained by contacting Heidi Resnick, Ph.D., National Crime Victims Research and Treatment Center, Department of Psychiatry and Behavioral Sciences, 171 Ashley Avenue, Charleston, SC 29425, USA. The Brief Trauma Interview, designed to briefly assess lifetime trauma exposure according to DSM-IV criteria in a clinically sensitive format, is also based on this measure. More information can be obtained by contacting Paula P. Schnurr, Ph.D., National Center for PTSD (116D), VA Medical and Regional Office Center, White River Junction, VT 05009, USA; (tel) 802-296-5132; (fax) 802-296-5135; (e-mail) paula.p.schnurr@dartmouth.edu.

Trauma History Questionnaire (THQ)

The THQ is a 24-item measure that assesses history of exposure to traumatic events. It is based on the High Magnitude Stressor Questionnaire from the DSM-IV field trials. It is currently considered an experimental instrument and thus the authors require a data sharing agreement. Preliminary psychometrics are available in a paper by Green (1996). The scale is available from Bonnie L. Green, Ph.D., Department of Psychiatry, Georgetown University, 611 Kober Cogan Hall, Washington, DC 20007, USA; (tel) 202-687-4812; (fax) 202-687-6658.

Traumatic Experiences Inventory (TEI)

The TEI is a 38-item self-report measure of the existence, intensity, and duration of symptoms that develop in response to a crime or natural disaster in which lives were lost. The factors include avoidance, reexperiencing, increased arousal, and victimization. More information on the scale is available in Sprang (1997).

Traumatic Stress Institute Belief Scale (TSI)

The TSI is designed to measure disrupted cognitive schemas that reflect trauma-sensitive needs. It was developed by Laurie Anne Pearlman, Ph.D., at the Traumatic Stress Institute/ Center for Adult & Adolescent Psychotherapy LLC, 22 Morgan Farms Drive, South Windsor, CT 06074, USA; (tel) 860-644-2541; (fax) 860-644-6891; (e-mail) laurie.pearlman@snet.net. A published version of the TSI is in development with Western Psychological Services, 12031 Wilshire Boulevard, Los Angeles, CA 90025, USA; (tel) 800-648-8857; (fax) 310-478-7838; (e-mail) custsvc@wpspublish.com; (website) www.wpspublish.com.

Traumatic Stress Schedule (TSS)

The TSS is a measure designed to detect the occurrence and impact of exposure to traumatic events. Respondents are asked about the occurrence of six types of events, as well as any additional events, and they are queried about the impact of these events on their life. The scale is reprinted in a paper by Norris (1990). More information is available from Fran Norris, Ph.D., Department of Psychology, Georgia State University, University Plaza, Atlanta, GA 30303, USA; (tel) 404-651-1607; (fax) 404-651-1391; (e-mail) fnorris@gsu.edu.

World Assumption Scale (WAS)

The WAS is a 32-item self-report measure of assumptions that are presumed to be impacted by exposure to traumatic events. The WAS taps into assumptions about the benevolence of the world, the meaning of the world, and self-worth. The measure is reprinted in a paper by Janoff-Bulman (1996) and some psychometric data are also available. More information about the scale is available from Ronnie Janoff-Bulman, Ph.D., Department of Psychology, University of Massachusetts, Amherst, MA 01003, USA; (tel) 413-545-0264; (fax) 413-545-0996; (e-mail) janbul@psych.umass.edu.

REFERENCES

- Allen, J. G., Coyne, L., & Huntoon, J. (1998). Trauma pervasively elevates Brief Symptom Inventory profiles in inpatient women. *Psychological Reports*, 83, 499-513.
- Andrykowski, M. A., Cordova, M. J., Studts, J. L., & Miller, T. W. (1998). Posttraumatic stress disorder after treatment for breast cancer: Prevalence of diagnosis and use of the PTSD Checklist-Civilian Version (PCL-C) as a screening instrument. *Journal of Consulting and Clinical Psychology*, 66, 586-590.
- Asmundson, G. J. G., Cox, B. J., Larsen, D. K., Frombach, I. K., & Norton, G. R. (1999). Psychometric properties of the Accident Fear Questionnaire: An analysis based on motor vehicle accident survivors in a rehabilitation setting. *Rehabilitation Psychology*, 44, 373–387.
- Astin, M. C., Lawrence, K. J., & Foy, D. W. (1993). Posttraumatic stress disorder among battered women: Risk and resiliency factors. Violence and Victims, 8, 17–29.
- Blake, D. D., Weathers, F. W., Nagy, L. M., Kaloupek, D. G., Klauminzer, G., Charney, D., & Keane, T. M. (1990). A clinician rating scale for assessing current and lifetime PTSD: The CAPS-1. *The Behavior Therapist*, 13, 187– 188.
- Blanchard, E. B., Jones-Alexander, J., Buckley, T. C., & Forneris, C. A. (1996). Psychometric properties of the PTSD Checklist (PCL). Behaviour Research and Therapy, 34, 669–673.
- Breslau, N., Kessler, R. C., & Peterson, E. L. (1998). PTSD assessment with a structured interview: Reliability and concordance with a standardized clinical interview. *International Journal of Methods in Psychiatric Research*, 7, 121–127.
- Briere, J., & Runtz, M. (1989). The Trauma Symptom Checklist (TSC-33): Early data on a new scale. Journal of Interpersonal Violence, 4, 151-163.
- Brom, D., Kleber, R. J., & Defares, P. B. (1989). Brief psychotherapy for posttraumatic stress disorders. Journal of Consulting and Clinical Psychology, 57, 607-612.
- Bryant, R. A., Harvey, A. G., Dang, S. T., & Sackville, T. (1998). Assessing acute stress disorder: Psychometric properties of a structured clinical interview. *Psychological Assessment*, 10, 215–220.
- Bryant, R. A., Moulds, M. L., & Guthrie, R. M. (2000). Acute Stress Disorder Scale: A self-report measure of acute stress disorder. *Psychological Assessment*, 12, 61–68.
- Cardeña, E., Koopman, C., Classen, C., Waelde, L. C., & Spiegel, D. (2000). Psychometric properties of the Stanford Acute Stress Reaction Questionnaire: A valid and reliable measure of acute stress. *Journal of Traumatic Stress*, 13, 719-734.
- Carlier, I. V. E., Lamberts, R. D., Van Uchelen, A. J., & Gersons, B. P. R. (1998). Clinical utility of a brief diagnostic test for posttraumatic stress disorder. *Psychosomatic Medicine*, 60, 42–47.
- Caplson, E. B., Putnam, F. W., Ross, C. A., Anderson, G., Clark, P., Torem, M., Coons, P., Bowman, E., Chu, J. A., Dill, D., Loewenstein, R. J., & Braun, B. G. (1991). Factor analysis of the Dissociative Experiences Scale: A multicenter study. In B. G. Braun & E. B. Carlson (Eds.), Proceedings of the Eighth International Conference on Multiple Personality and Dissociative States. Chicago: Rush.
- Dalton, J. E., Tom, A., Rosenblum, M. L., Garte, S. H., & Aubuchon, I. N. (1989). Faking on the Mississippi Scale for Combat-Related Posttraumatic Stress Disorder. *Psychological Assessment*, 1, 56–57.
- Davidson, J. R. T., & Colket, J. T. (1997). The eight-item Treatment-Outcome Post-Traumatic Stress Disorder Scale: A brief measure to assess treatment outcome in post-traumatic stress disorder. *International Clinical Psychophar*macology, 12, 41–45.
- Davidson, J. R., Kudler, H. S., Saunders, W. B., Erickson, L., Smith, R. D., Stein, R. M., Lipper, S., Hammett, E. B., Mahorney, S. L., & Cavenar, J. O. (1993). Predicting response to amitriptyline in posttraumatic stress disorder. *American Journal of Psychiatry*, 150, 1024–1029.

- Dutton, M. A., Perrin, S., Chrestman, K., Halle, P., & Burghardt, K. (1991, August). Posttraumatic stress disorder in battered women: Concurrent validity. Paper presented at the 99th Annual Convention of the American Psychological Association, San Francisco.
- Falsetti, S., Resnick, H., Kilpatrick, D., & Freedy, J. (1994). A review of the Potential Stressful Events Interview. *The Behavior Therapist*, 17, 66–67.
- Falsetti, S. A., Resnick, H. S., Resick, P. A., & Kilpatrick, D. G. (1993). The Modified PTSD Symptom Scale: A brief self-report measure of posttraumatic stress disorder. *Behavioral Assessment Review*, 16, 161–162.
- Fischer, D., & Elnitsky, S. (1990). A factor analytic study of two scales measuring dissociation. American Journal of Clinical Hypnosis, 32, 201–207.
- Foa, E. B., Riggs, D. S., Dancu, C. V., & Rothbaum, B. O. (1993). Reliability and validity of a brief instrument for assessing posttraumatic stress disorder. *Journal of Traumatic Stress*, 6, 459–473.
- Foa, E. B., Rothbaum, B. O., Riggs, D. S., & Murdock, T. B. (1991). Treatment of posttraumatic stress disorder in rape victims: A comparison between cognitive-behavioral procedures and counseling. *Journal of Consulting and Clinical Psychology*, 59, 715–723.
- Fontana, A., & Rosenheck, R. (1994). A short form of the Mississippi Scale for measuring change in combat-related PTSD. Journal of Traumatic Stress, 7, 407–414.
- Foy, D. W., Sipprelle, R. C., Rueger, D. B., & Carroll, E. M. (1984). Etiology of posttraumatic stress disorder in Vietnam veterans: Analysis of premilitary, military, and combat exposure influences. *Journal of Consulting and Clinical Psychology*, 52, 79–87.
- Foy, D. W., Wood, J. L., King, D. W., King, L. A., & Resnick, H. S. (1997). Los Angeles Symptom, Checklist: Psychometric evidence with an adolescent sample. Assessment, 4, 377–384.
- Frischholz, E. J., Braun, B. G., Sachs, R. G., Hopkins, L., Shaeffer, D. M., Lewis, J., Leavitt, F., Pasquotto, M. A., & Schwartz, D. R. (1990). The Dissociative Experiences Scale: Further replication and validation. *Dissociation*, 3, 151–153.
- Frueh, B. C., Turner, S. M., Beidel, D. C., Mirabella, R. F., & Jones, W. J. (1996). Trauma Management Therapy: A preliminary evaluation of a multicomponent behavioral treatment for chronic combat-related PTSD. *Behaviour Research and Therapy*, 34, 533-543.
- Gold, J. W., & Cardeña, E. (1998). Convergent validity of three posttraumatic symptoms inventories among adult sexual abuse survivors. *Journal of Traumatic Stress*, 11, 173–180.
- Green, B. L. (1996). Psychometric review of Trauma History Questionnaire (Self-Report). In B. H. Stamm (Ed.), Measurement of stress, trauma and adaptation (pp. 366–388). Lutherville, MD: Sidran Press.
- Green, B. L., Goodman, L. A., Krupnick, J. L., Corcoran, C. B., Petty, R. M., Stockton, P., & Stern, N. M. (2000). Outcomes of single versus multiple trauma exposure in a screening sample. *Journal of Traumatic Stress*, 13, 271–286.
- Hammarberg, M. (1992). Penn Inventory for Posttraumatic Stress Disorder: Psychometric properties. Psychological Assessment, 4, 67–76.
- Hammarberg, M., & Silver, S. M. (1994). Outcome of treatment for post-traumatic stress disorder in a primary care unit serving Vietnam veterans. *Journal of Traumatic Stress*, 7, 195–216.
- Hendrix, C. C., Jurich, A. P., & Schumm, W. R. (1994). Validation of the Impact of Event Scale on a sample of American Vietnam veterans. *Psychological Reports*, 75, 321-322.
- Herman, D. S., Weathers, F. W., Litz, B. T., & Keane, T. M. (1996). Psychometric properties of the embedded and stand-alone versions of the MMPI-2 Keane PTSD scale. Assessment, 3, 437-442.
- Hodgkinson, P., & Joseph, S. (1995). Factor analysis of the Impact of Event Scale with female bank staff following an armed raid. *Personality and Individual Differences*, 19, 773–775.
- Horowitz, M. (1976). Stress response syndromes. New York: Jason Aronson.
- Horowitz, M. (1982). Stress response symptoms and their treatment. In L. Goldberger & S. Breznitz (Eds.), Handbook of stress: Theoretical and clinical aspects (pp. 711-732). New York: Free Press.
- Housekamp, B. M., & Foy, D. W. (1991). The assessment of posttraumatic stress disorder in battered women. Journal of Interpersonal Violence, 6, 368–376.
- Hovens, J. E., van der Ploeg, H. M., Klaarenbeek, M. T. A., Schreuder, J. N., & Rivero, V. V. (1994a). The assessment of posttraumatic stress disorder with the Clinician Administered PTSD Scale: Dutch results. *Journal of Clinical Psychology*, 50, 325–340.
- Hovens, J. E., van der Ploeg, H. M., Klaarenbeek, M. T. A., Schreuder, J. N., & Rivero, V. V. (1994b). The development of the Self-Rating Inventory for Posttraumatic Stress Disorder. Acta Psychiatrica Scandinavica, 90, 172–183.
- Hyer, L., Summers, M. N., Boyd, S., Litaker, M., & Boudewyns, P. (1996). Assessment of older combat veterans with the Clinician-Administered PTSD Scale. *Journal of Traumatic Stress*, 9, 587–593.
- Janoff-Bulman, R. (1996). Psychometric review of World Assumption Scale. In B. H. Stamm (Ed.), Measurement of stress, trauma and adaptation (pp. 440-442). Lutherville, MD: Sidran Press.

- Joseph, S. (2000). Psychometric evaluation of Horowitz's Impact of Event Scale: A review. Journal of Traumatic Stress, 13, 101-113.
- Joseph, S., Yule, W., Williams, R., & Hodgkinson, P. (1993). The Herald of Free Enterprise disaster: Measuring posttraumatic symptoms 30 months on. British Journal of Clinical Psychology, 32, 327-331.
- Kenderdine, S. K., Phillips, E. J., & Scurfield, R. M. (1992). Comparison of the MMPI-PTSD subscale with PTSD and substance abuse patient populations. *Journal of Clinical Psychology*, 48, 136–139.
- King, D. W., Leskin, G. A., King, L. A., & Weathers, F. W. (1998). Confirmatory factor analysis of the Clinician-Administered PTSD Scale: Evidence for the dimensionality of posttraumatic stress disorder. *Psychological Assessment*, 10, 90–96.
- King, L. A., & King, D. W. (1994). Latent structure of the Mississippi Scale for Combat-Related Post-Traumatic Stress Disorder: Exploratory and higher-order confirmatory factor analyses. Assessment, 1, 275–291.
- Koretzky, M. B., & Peck, A. H. (1990). Validation and cross-validation of the PTSD subscale of the MMPI with civilian trauma victims. *Journal of Clinical Psychology*, 46, 296–300.
- Koss, M. P., Gidycz, C. A., & Wisniewski, N. (1987). The scope of rape: Incidence and prevalence of sexual aggression and victimization in a national sample of higher education students. *Journal of Consulting and Clinical Psychology*, 55, 162-170.
- Lauterbach, D., Vrana, S., King, D. W., & King, L. A. (1997). Psychometric properties of the civilian version of the Mississippi PTSD Scale. *Journal of Traumatic Stress*, 10, 499–513.
- Lawrence, K. J. (1992). Assessment of posttraumatic stress disorder among survivors of ritual abuse. Unpublished doctoral dissertation, Fuller Theological Seminary, Pasadena, CA.
- Leskin, G., & Foy, D. W. (1993, October). Development and psychometric validation of the LA Symptom Checklist. In L. A. King (Chair), Assessment of PTSD by self-report measure: Implications for construct validity. Symposium presented at the annual meeting of the International Society for Traumatic Stress Studies, San Antonio, TX.
- Litz, B. T., Orsillo, S. M., Kaloupek, D., & Weathers, F. (2000). Emotional-processing in posttraumatic stress disorder. Journal of Abnormal Psychology, 109, 26–39.
- Litz, B. T., Penk, W. E., Walsh, S., Hyer, L., Blake, D. D., Marx, B., Keane, T. M., & Bitman, D. (1991). Similarities and differences between MMPI and MMPI-2 applications to the assessment of posttraumatic stress disorder. *Journal of Personality Assessment*, 57, 238–253.
- Lund, M., Foy, D. W., Sipprelle, R. C., & Strachan, A. M. (1984). The Combat Exposure Scale: A systematic assessment of trauma in the Vietnam war. *Journal of Clinical Psychology*, 40, 1323–1328.
- Lyons, J. A., Caddell, J. M., Pittman, R. L., Rawls, R., & Perrin, S. (1994). The potential for faking on the Mississippi Scale for Combat-Related PTSD. Journal of Traumatic Stress, 7, 441–445.
- Malekzai, A. S. B., Niazi, J. M., Paige, S. R., Hendricks, S. E., Fitzpatrick, D., Leuschen, M. P., & Millimet, C. R. (1996). Modification of CAPS-1 for diagnosis of PTSD in Afghan refugees. *Journal of Traumatic Stress*, 9, 891–898.
- Manne, S. L., Du Hamel, K., Gallelli, K., Sorgen, K., & Redd, W. H. (1998). Posttraumatic stress disorder among mothers of pediatric cancer survivors: Diagnosis, comorbidity, and utility of the PTSD Checklist as a screening instrument. Journal of Pediatric Psychology, 23, 357–366.
- Marmar, C. R., Weiss, D. S., & Metzler, T. (1997). The Peritraumatic Dissociative Experiences Questionnaire. In J. P. Wilson & T. M. Keane (Eds.), Assessing psychological trauma and PTSD (pp. 412–428). New York: Guilford.
- McDonald, A. S. (1997). Factor structure of the Impact of Event Scale in a non-clinical sample. Personality and Individual Differences, 23, 419-424.
- McFall, M. E., Smith, D. E., Mackay, P. W., & Tarver, D. J. (1990). Reliability and validity of Mississippi Scale for Combat-Related Posttraumatic Stress Disorder. *Psychological Assessment*, 2, 114–121.
- McFall, M. E., Smith, D. E., Roszell, D. K., Tarver, D. J., & Malas, K. L. (1990). Convergent validity of measures of PTSD in Vietnam combat veterans. American Journal of Psychiatry, 147, 645-648.
- Meltzer-Brody, S., Churchill, E., & Davidson, J. R. T. (1999). Derivation of the SPAN, a brief diagnostic screening test for post-traumatic stress disorder. *Psychiatry Research*, 88, 63-70.
- Moody, D. R., & Kish, G. B. (1989). Clinical meaning of the Keane PTSD scale. Journal of Clinical Psychology, 45, 542–546.
- Munley, P. H., Bains, D. S., Bloem, W. D., & Busby, R. M. (1995). Post-traumatic stress disorder and the MMPI-2. Journal of Traumatic Stress, 8, 171-178.
- Neal, L. A., Busuttil, W., Herapath, R., & Strike, P. W. (1994). Development and validation of the computerized Clinician Administered Posttraumatic Stress Disorder Scale-1-Revised. *Psychological Medicine*, 24, 701–706.
- Neal, L. A., Busuttil, W., Rollins, J., Herepath, R., Strike, P., & Turnbull, G. (1994). Convergent validity of measures of post-traumatic stress disorder in a mixed military and civilian population. *Journal of Traumatic Stress*, 7, 447– 455.
- Norris, F. H. (1990). Screening for traumatic stress: A scale for use in the general population. Journal of Applied Social Psychology, 20, 1704–1718.

- Norris, F. H., & Perilla, J. L. (1996). The Revised Civilian Mississippi Scale for PTSD: Reliability, validity, and crosslanguage stability. Journal of Traumatic Stress, 9, 285–298.
- Pava, W. (1993). {Administration of LASC}. Unpublished data.
- Perry, S., Difede, J., Musngi, G., Frances, A. J., & Jacobsberg, L. (1992). Predictors of posttraumatic stress disorder after burn injury. American Journal of Psychiatry, 149, 931-935.
- Pitblado, C. B., & Sanders, B. (1991). Reliability and short-term stability of scores on the Dissociative Experiences Scale. In B. G. Braun & E. B. Carlson (Eds.), Proceedings of the Eighth International Conference on Multiple Personality and Dissociative States. Chicago: Rush.
- Resnick, H. S., Foy, D. W., Donahoe, C. P., & Miller, E. N. (1989). Antisocial behavior and posttraumatic stress disorder in Vietnam veterans. *Journal of Clinical Psychology*, 45, 860–866.
- Ross, C. A., Joshi, S., & Currie, R. (1991). Dissociative experiences in the general population: A factor analysis. Hospital and Community Psychiatry, 42, 297–301.
- Rowan, A. B., Foy, D. W., Rodriguez, N., & Ryan, S. (1994). Posttraumatic stress disorder in adults sexually abused as children. International Journal of Child Abuse and Neglect, 18, 51-61.
- Ryan, S. (1992). Psychometric analysis of the Sexual Abuse Exposure Questionnaire. Unpublished doctoral dissertation, Fuller Theological Seminary, Pasadena, CA.
- Saunders, B. E., Arata, C. M., & Kilpatrick, D. G. (1990). Development of a crime-related post-traumatic stress disorder scale for women within the Symptom Checklist-90-Revised. Journal of Traumatic Stress, 3, 439–448.
- Schwarzwald, J., Solomon, Z., Weisenberg, M., & Mikulincer, M. (1987). Validation of the Impact of Event Scale for psychological sequelae of combat. *Journal of Consulting and Clinical Psychology*, 55, 251–256.
- Scotti, J. R., Sturges, L. V., & Lyons, J. A. (1996). The Keane PTSD Scale extracted from the MMPI: Sensitivity and specificity with Vietnam veterans. *Journal of Traumatic Stress*, 9, 643–650.
- Shalev, A. Y., Freedman, S., Peri, T., Brandes, D., & Sahar, T. (1997). Predicting PTSD in trauma survivors: Prospective evaluation of self-report and clinician-administered instruments. *British Journal of Psychiatry*, 170, 558-564.
- Shalev, A. Y., Peri, T., Canetti, L., & Schreiber, S. (1996). Predictors of PTSD in injured trauma survivors: A prospective study. American Journal of Psychiatry, 153, 219–225.
- Solomon, Z., Benbenishty, R., Neria, Y., Abramowitz, M., Ginzburg, K., & Ohry, A. (1993). Assessment of PTSD: Validation of the revised PTSD Inventory. Israel Journal of Psychiatry and Related Sciences, 30, 110-115.
- Sprang, G. (1997). The Traumatic Experiences Inventory (TEI): A test of psychometric properties. Journal of Psychopathology and Behavioral Assessment, 19, 257-271.
- Thompson, J. A., Charlton, P. F. C., Kerry, R., & Lee, D. (1995). An open trial of exposure therapy based on deconditioning for post-traumatic stress disorder. British Journal of Clinical Psychology, 34, 407-416.
- van Ijzendoorn, M. H., & Schuengel, C. (1996). The measurement of dissociation in normal and clinical populations: Meta-analytic validation of the Dissociative Experiences Scale (DES). Clinical Psychology Review, 16, 365-382.
- Vrenen, D. L., Gudanowski, D. M., King, L. A., & King, D. W. (1995). The civilian version of the Mississippi PTSD Scale: A psychometric evaluation. *Journal of Traumatic Stress*, 8, 91–109.
- Walker, E. A., Katon, W. J., Neraas, K., Jemelka, R. P., & Massoth, D. (1992). Dissociation in women with chronic pelvic pain. American Journal of Psychiatry, 149, 534–537.
- Waller, N. G. (1995). The Dissociative Experiences Scale. In J. C. Conoley & J. C. Impara (Eds.), Twelfth mental measurements yearbook (pp. 317–318). Lincoln, NE: Buros Institute of Mental Measurement.
- Waller, N. G., Putnam, F. W., & Carlson, E. B. (1996). Types of dissociation and dissociative types: A taxometric analysis of dissociative experiences. *Psychological Methods*, 1, 300–321.
- Waller, N. G., & Ross, C. A. (1997). The prevalence and biometric structure of pathological dissociation in the general population: Taxometric and behavior genetic findings. *Journal of Abnormal Psychology*, 106, 499–510.
- Watson, C. G., Juba, M. P., Manifold, V., Kucala, T., & Anderson, P. E. D. (1991). The PTSD Interview: Rationale, description, reliability, and concurrent validity of a DSM-III-based technique. *Journal of Clinical Psychology*, 47, 179–188.
- Watson, C. G., Plemel, D., DeMotts, J., Howard, M. T., Tuorila, J., Moog, R., Thomas, D., & Anderson, D. (1994). A comparison of four PTSD measures' convergent validities in Vietnam veterans. *Journal of Traumatic Stress*, 7, 75-82.
- Weathers, F. W., & Litz, B. T. (1994). Psychometric properties of the Clinician Administered PTSD Scale, CAPS-1. PTSD Research Quarterly, 5, 2–6.
- Weathers, F. W., Litz, B. T., Keane, T. M., Herman, D. S., Steinberg, H. R., Huska, J. A., & Kraemer, H. C. (1996). The utility of the SCL-90-R for the diagnosis of war-zone related posttraumatic stress disorder. *International Society* for Traumatic Stress Studies, 9, 111–128.
- Weathers, F. W., Ruscio, A. M., & Keane, T. M. (1999). Psychometric properties of nine scoring rules for the Clinician Administered Posttraumatic Stress Disorder Scale. *Psychological Assessment*, 11, 124–133.

MEASURES FOR ACUTE STRESS DISORDER AND PTSD | 307

- Weiss, D. S., Horowitz, M. J., & Wilner, N. (1984). The Stress Response Rating Scale: A clinician's measure for rating the response to serious life-events. *British Journal of Clinical Psychology*, 23, 202–215.
- Weiss, D. S., & Marmar, C. R. (1997). The Impact of Event Scale-Revised. In J. P. Wilson & T. M. Keane (Eds.), Assessing psychological trauma and PTSD (pp. 98–135). New York: Guilford.
- Wolfe, J., & Kimerling, R. (1997). Gender issues in the assessment of posttraumatic stress disorder. In J. P. Wilson & T. M. Keane (Eds.), Assessing psychological trauma and PTSD (pp. 192–238). New York: Guilford.
- Zilberg, N. J., Weiss, D. S., & Horowitz, M. J. (1982). Impact of Event Scale: A cross-validation study and some empirical evidence supporting a conceptual model of stress response syndromes. *Journal of Consulting and Clinical Psychology*, 50, 407–414.
- Zlotnick, C., Davidson, J., Shea, M. T., & Pearlstein, T. (1996). Validation of the Davidson Trauma Scale in a sample of survivors of childhood sexual abuse. *Journal of Nervous and Mental Disease*, 184, 255-257.
- Zlotnick, C., & Pearlstein, T. (1997). Validation of the Structured Interview for Disorders of Extreme Stress. Comprehensive Psychiatry, 38, 243-247.
- Zlotnick, C., Shea, M., Zakriski, A., Costello, E., Begin, A., Pearlstein, T., & Simpson, E. (1995). Stressors and close relationships during childhood and dissociative experiences in survivors of sexual abuse among inpatient psychiatric women. *Comprehensive Psychiatry*, 36, 207–212.

÷